Texas School Survey of Substance Use Methods Report 2006

For the Texas Department of State Health Services

Prepared by

James A. Dyer, Ph.D. Melissa Tackett-Gibson, Ph.D. Ashweeta Patnaik

Public Policy Research Institute
Texas A&M University
College Station, TX 77843-4476

Introduction

The Public Policy Research Institute (PPRI), in conjunction with the Department of State Health Services, conducted the ninth statewide survey of drug and alcohol use among Texas elementary and secondary students in the Spring of 2006. Originally implemented in 1988 as a component of a larger survey assessing substance use among the state's general population, the school survey has since become an ongoing, independent project. District surveys are offered every year with a statewide survey conducted every two years. The 2006 effort provides follow-up data reflecting changes over the past fourteen years in grades four through twelve.

The *Texas School Survey* project has two primary objectives. First, it serves to inform state and local policy-makers about the extent and nature of the substance use problem in Texas schools. Second, the statewide survey provides a standard of comparison for districts conducting local assessments of drug and alcohol use.

The purpose of this document is to describe the methodology used to administer the 2006 *Texas School Survey of Substance Use*. Following a brief introduction to the survey instrument itself, attention is then focused on sample selection and survey administration procedures. Next, methods for data processing and quality control are described and the report concludes with a review of standard error estimates.

Survey Instrument

Two versions of the 2006 Texas School Survey of Substance Use were developed and administered. The first was a six-page questionnaire designed for students in grades seven through twelve. The second was a simplified three-page instrument created for students in grades four through six. The elementary survey differs from the secondary survey in that it has simplified language and some complex questions were omitted. Elementary students were asked about only four types of substances including tobacco (cigarettes, snuff, and chewing tobacco), alcohol (beer, wine, wine coolers, and liquor), inhalants, and marijuana. Secondary students were asked about the same substances, as well as a broader range of illicit drugs including powdered cocaine, crack, hallucinogens, uppers, downers, steroids, ecstasy, Rohypnol, DXM, and heroin. Other sets of questions on both the elementary and secondary instruments were designed to assess behavioral correlates of substance use and students' perceptions of support available to help them cope with substance-related problems.

The questionnaire was in a format that could be scanned optically, similar to that used for standardized testing. It was designed for anonymous self-administration by students with the aid of a staff member to pass out the survey, read a common set of instructions, monitor the class during survey administration, and collect the instruments after they are completed. The survey instruments are included in Appendix A.

Survey Content

The Texas School Survey of Substance Use content was updated in 2006.

The *Texas School Survey of Substance Use* content was updated in slightly in 2006. Changes were made in the responses to participation in extra-curricular school activities to make a better distinction between those activities that were directly school related activities and those that were not(Q12). There were wording changes in the responses to binge the binge drinking question to

make it consistent with reporting requirements (Q26). Responses to questions on where student had received information about alcohol and drugs in school and out of school were changed to reflect current options(Q37,Q38). A question about being tested for drugs by school was added (Q40).

Survey Sample

The sample of students for the 2006 survey was designed to be a stratified, weighted, random sample of all public school students between the fourth and twelfth grades in the state. In order to make administration practical, students were selected using a multi-stage stratified sampling procedure. This involved sampling districts, schools within districts, and classrooms within districts. All students in a sampled classroom were asked to participate in the survey.

The 2006 sample design was a modification of the design used since 2000. The number of strata used was increased so that it would be possible to provide estimates for each of the eight state planning regions.

Selection of Districts

The primary analytic cluster was the school district since the approval needed to administer the survey had to be obtained at that level. First, the districts were stratified into the eight health planning regions and further divide each of these regions areas based on their status in relationship to metropolitan areas (metro status). Finally, within the strata defined by region and metro status the districts were further divided by size of district. Finally, all border districts were sample in order to minimize the error for comparisons between the border and non-border regions of the state. The strata are listed in Table 1 below.

The strata with the number of districts sampled from each strata was a function of the size of the enrollment represented in the strata and the number of districts. To minimize error in the sample, all large districts were sample. Districts with smaller numbers of enrollment contribute less error to the analysis as they involve a smaller proportion of students. In strata where only some districts were to be selected, districts were randomly selected. Where possible, additional districts were sampled in each strata in replace those not participating. In Table 1, the numbers of districts in the strata are listed along with the number of districts targeted. The total list of districts participating by strata is provided in Appendix B.

TABLE 1. Sample Strata Definitions and Target Sample

Adminstrative Region	Metro Status	<=10,000	<=5,000	>11,000- 28,000	>15,000	>28,000- 43,000	>43,000	>5,000	>5,000- 15,000	Grand Total
1	Metro Central City	3		1		1				5
	Metro Suburban Non-Metro		2						1	3
	Adjacent Non-Metro Non- Adjacent		7 5					1		8 5
7	Metro Central City	3		2		1	1			7
	Metro Suburban Non-Metro Adjacent		5 7		2				3	10 7
8	Metro Central City	3		2			3			8
	Metro Suburban		5						3	8
	Non-Metro Adjacent Non-Metro Non-		12							12
	Adjacent		4					1		5
11	Metro Central City	21		9		2	1			33
	Metro Suburban Non-Metro		2							2
	Adjacent		22					3		25
2/3	Metro Central City	3	_	5		3	4			15
	Metro Suburban Non-Metro Adjacent		7		2			3	3	12 6
	Non-Metro Non- Adjacent		3					3		6
4/5	Metro Central City	7		1						8
	Metro Suburban Non-Metro		3					4		3
	Adjacent Non-Metro Non- Adjacent		17					1		18
6/5	Metro Central City	3		1		1				5
	Metro Suburban Non-Metro		5		2				2	9
	Adjacent Non-Metro Non- Adjacent		3					17 2		20
9/10	Metro Central City	9		3		1	2			15
<i>5,</i> 10	Non-Metro Adjacent		5			'		1		6
	Non-Metro Non- Adjacent		13							13
Grand Total	. ,	52	133	24	6	9	11	32	12	279

Soliciting District Participation

Obtaining cooperation from those districts that were randomly selected for the state sample when the selected district did not plan to do a local survey was sometimes a problem. Yet, it was critical to get data from as many of the originally selected districts as possible. Some state sample districts that were initially hesitant were persuaded to cooperate by the use of incentives. The various incentives used included waiving participation and sampling fees, offering to discount the fees for participating the following year, paying all shipping costs, and discounting campus level analyses fees.

In order to allow further analysis of substance use among students living on the Texas-Mexico border, school districts along the border were strongly encouraged to participate in the 2006 *Texas School Survey*. The survey was offered free of charge to border districts, and data was collected from a broadly defined 28-county area. The border sample was designed to collect data from approximately 500,000 students. One-hundred and eight school districts from 28 counties were invited to participate in the 2006 survey.

Participation Rates

Sixty-three of the original 279 selected districts participated in the study. One-hundred and twenty-three districts were not able to participate; and most declined due to the lack of time and resources involved in survey administration. In lieu of the declining districts, an additional twenty districts were included as alternates for the final sample. With alternates, the sample included 203 districts. In total 81 districts participated in the 2006 survey.

The selected districts included 2.87 million elementary and secondary students and those that were included in the final sample included 1.35 million students. A total of 47 percent of the students in the original sample were in the final sampling frame.

Seventy-one of the 81 districts sampled submitted both elementary and secondary surveys. Ten districts, Beaumont, Boerne, College Station, Conroe, Cypress-Fairbanks, East Chambers, Eustace, Hooks, Nursery, and South Texas, did not administer surveys to elementary students. All of the districts surveyed middle school and high school students.

Allocation of Surveys among Districts

The state survey sample was designed to collect data from a minimum sample of about 5,555 students per grade, however, many districts chose to survey more than the minimum number of students specified in the state sampling plan. Some extremely small districts received somewhat more than a strict proportional allocation because, while the data was technically only needed from one or two students per grade, the survey was administered to the entire classroom. Similarly, in a few extremely large (urban) districts, fewer students were need for accuracy than would result from a true proportional allocation. All surveys submitted from a cooperating district were included in the sample. Accordingly, in the final analyses, the data were weighted to provide an accurate proportional allocation.

Thus, although we had estimated that the state sample would include approximately 50,000 students, it actually included 74,442 elementary students and 141,905 secondary students (See Table 2). This significantly improves the accuracy of estimates.

TABLE 2. Number of Surveys Included in State Sample

Allocation of Surveys among Classrooms and Campuses

Once the number of surveys to be administered in each district was established, the next step was to determine the number of classrooms to be surveyed per grade. This was achieved by dividing the number of questionnaires per grade (ascertained for each district using proportional population calculations) by the average number of students per class---20 for grades four through six, 22 for grades seven through twelve. The result of this computation indicated the total number of classes to be surveyed. These classes were selected so that as many different campuses as possible were in the final sample. Ideally, the classrooms surveyed were evenly distributed across all campuses in the district. If there were more campuses containing a given grade than classrooms needed, then a simple random selection procedure was used to determine which campuses would be sampled. In general, once a campus was selected, all relevant grades at that campus were surveyed. Therefore, campus selection was not independent between grades.

TABLE 2. Survey Distribution by Grade

	Grade	Number of Usable Surveys	Percentage
Elementary	4^{th}	21,560	29%
•	5 th	26,153	35%
	6^{th}	26,729	36%
		74,442	100%
Secondary	7^{th}	29,646	21%
•	8^{th}	25,484	18%
	9 th	26,966	19%
	10^{th}	21,455	15%
	11^{th}	20,971	15%
	12^{th}	17,383	12%
		141,905	100%

Selection of Classrooms within Campuses

After the total number of classrooms to be surveyed in each grade at each campus was determined, it was necessary to identify specific classrooms. This selection procedure was performed by campus personnel based on a set of guidelines provided by PPRI (illustrated in Appendix C). Members of campus staff were asked to make a list by grade (according to teacher's last name or some other convenient method) of all classes held during a selected class period. They were then instructed to use a random number table to select the exact classes to survey in each grade.

Other Sampling Considerations

Some school districts sampled all students in all or some of the grades. In these districts, the methodology outlined above did not apply to the grades sampled at 100 percent. In Houston the district used a list of all students from which to conduct a random sample of the students. Therefore, there are no campuses and classrooms sampled in these districts.

Survey Administration Procedures

Districts selected for inclusion in the state sample were notified about the project via letter and were sent a descriptive brochure, illustrated in Appendix D. State sample districts that planned to administer a local drug and alcohol survey had virtually no procedural changes resulting from their involvement in the statewide project. In those districts that surveyed grades four through twelve, sufficient data was collected from all relevant campuses to meet the data collection needs of the statewide survey. These districts benefited from their inclusion in the state survey project because they were not charged for the surveys that became part of the state database. The larger number of surveys from these districts was weighted down so that their contribution to the final sample was in correct proportion.

In those instances where state sample districts were collecting local data for an incomplete combination of grades, or where they were not conducting local surveys at all, the campus and classroom selection procedures described above were applied. Arrangements for giving the survey were established on an individual basis with these districts. Since those not doing local surveys did not stand to gain directly from having the survey administered in their district, an effort was made to be as accommodating as possible. PPRI was able to arrange survey administration in the selected schools and classes by school personnel.

In some larger districts, such as Houston and Austin, the district uses the computer to draw a random sample of all students. On each campus where the students are located, the students are requested to go to a specified room where the survey is conducted. Once in the room, the survey is conducted, as it would be in a classroom in the other districts.

Relevant personnel in the selected districts and campuses were provided with complete instructions and materials necessary to administer the survey (see Appendix E). Classrooms were selected randomly by PPRI based on information from a computer printout from the district or Campus Information Form. Teachers in selected classrooms were given a script to read so that all students would receive a standardized set of instructions. Teachers were also asked to complete a Classroom Identification Form that provided data on the number of students that should have taken the survey but were absent, and the number that was present but failed to complete the survey. This information was useful for computing error estimates. After the surveys were administered in each classroom, they were sealed in an envelope along with the Classroom Identification Form. The envelopes from all participating classrooms were collected and returned to PPRI.

Data Entry and Analyses

As noted earlier, the format of the survey instruments enabled them to be scanned optically. Upon receipt at PPRI, the instruments were logged in, coded, and scanned by staff or trained personnel.

Exaggerated Responses

Because the *Texas School Survey* data is based entirely upon respondents' description of their own behavior, it is inevitable that some students will under- or over-report their use of drugs or alcohol, and to the extent possible PPRI attempted to identify and eliminate data from those respondents. Two checks were incorporated into the data analysis program to identify exaggerators. First, both elementary and secondary students were asked about their use of a false drug call "Shimeron." Data from students claiming to have used this substance were considered suspect and dropped from the analyses.

Second, checks were run to identify any students claiming extremely high levels of drug and alcohol use. Unbelievable high substance use for elementary students was defined as the use of five or more substances, 11 or more times in the past school year or over a lifetime. Secondary students were defined as exaggerators based on the following criteria: (1) students reported that they had five or more drinks of two or more beverages every day; (2) students reported that they had consumed three or more alcoholic beverages every day; or (3) students reported that they used four or more drugs (other than cigarettes, alcohol, or steroids) eleven or more times in the past month. As in those cases where students reported using "Shimeron," data from students reporting exaggerated use were also dropped from the analyses. Less than two percent (1.23%) of the total elementary sample exaggerated. The percentage of secondary school students who exaggerated (3.83%) was more than twice that of elementary students.

Unreported Grade Levels

When students failed to report their grade level, it was impossible to determine unequivocally in which grade these students' data should be analyzed. Approximately, 1% of the elementary and secondary surveys were missing grade information. When a grade level was missing, an estimate of the grade was made based on the students' age and the data were retained. Table 6 identifies the range of students' ages and the corresponding grade levels that were assigned. If both grade and age were missing, the data were dropped from the analyses.

TABLE 6. Age-Based Grade Assignments.

Age	Elementary Level	Age	Secondary Level
9	4 th Grade	12	7 th Grade
10	5 th Grade	13	8 th Grade
11	6 th Grade	14	9 th Grade
		15	10 th Grade
		16	11 th Grade
		17 or older	12 th Grade

Quality Control Measures

To ensure the quality of the statewide survey data, a number of internal checks were put into place to guide survey processing. First, a quality control analyst oversaw the implementation of all pre- and post-analysis quality control procedures. As the following paragraphs describe, many aspects of PPRI's plan for quality control were embedded in automated procedures. However, there is no replacement for human oversight. The quality control analyst monitored and tracked the processing of each district's surveys from the initial mailing through the production of the final state report. Responsibilities included ensuring that surveys were properly coded and scanned and checking for anomalies in the final table of results.

In addition to the safeguards resulting from careful project oversight, there were also a number of procedural checks against error. For example, there was a possibility, however remote, that after the bindings of a set of survey instruments were cut, the instruments could be dropped or otherwise placed out of order. If this occurred, it is conceivable that some pages of data could have been read into the incorrect computer record. To resolve this problem, each instrument used in the 2006 survey was printed with a five-digit "litho-code" number. With this coding process, every page of a given instrument is printed with the same scannable number, but a unique number is assigned to every instrument. By using the litho-code, when each page of an instrument is scanned it will automatically be read into the correct computer record. In this way, even if the pages from different instruments were shuffled together and read randomly, all data derived from the same instrument would automatically be read to the same data record.

Litho-coding also enabled PPRI to confirm that data from every survey instrument read was associated with the correct district. Survey instruments were mailed to participating districts in consecutive order. By recording the beginning and ending instrument numbers going to each district, PPRI was able to check the litho-codes scanned for a given district. In this way, any stacks of data that could potentially have been inadvertently mislabeled could be easily identified.

Programming checks were also incorporated into the data analysis program by cross-analysis. That is, the same data was run in several different ways using existing programs, and program outputs were then compared for consistency. Confidence is high that these quality control features will ensure valid and reliable survey findings.

Weights, Standard Errors, and Confidence Intervals

Weights were applied to each case based on the strata, district, and campus. The weights were applied so that the aggregation of students in each campus, district, and strata reflected their proportions in the actual district, campus, and strata populations. The formulae use to determine these weights are presented in Appendix F.

Standard errors and confidence intervals were estimated for each grade and the aggregation. The formulae used are presented in Appendix G. The table of standard errors and confidence intervals for past year and lifetime use of substances by grades are presented in Appendix H.

Conclusion

The *Texas School Survey* has become a valuable policy tool for both state and local educators and policy-makers. The survey, performed every two years, provides timely and relevant information about current drug and alcohol use patterns among young people enrolled in the Texas' public schools. Furthermore, longitudinal analysis can provide insight into changes in drug and alcohol prevalence over time. As was noted in the introduction, every state survey culminates in a Department of State Health Services publication providing an overview of findings to date. Data is also available for independent analysis by policy-makers and academicians.

APPENDIX A

Survey Instruments

TEXAS SCHOOL SURVEY OF SUBSTANCE USE

ELEMENTARY SURVEY INSTRUMENT 2006 VERSION

INTRODUCTION

Before you answer any questions in this survey, you need to know the following:

- This is a survey that measures tobacco, drug and alcohol use and other information relating to tobacco, alcohol, inhalant and drug and use;
- This survey is being given to as many as 300,000 students in the state of Texas;
- There is no way for anyone to know how you respond to the questions; and
- You can skip any question or not take the survey at all if you do not want to. There is no penalty if you skip questions or leave the survey blank.


If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol or drugs, counselors from the Worker's Assistant Program of Texas are available by calling this toll free number: 1-877-9-NO-DRUG (1-877-966-3784).


This research study has been approved and reviewed by the Institutional Review Board.—Human Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted through Ms. Angelia M. Raines, Director of Research Compliance, at (979) 458-4067 or email her at araines@vprmail.tamu.edu.

DIRECTIONS

- · DO NOT write your name anywhere on this booklet.
- · Use a NUMBER 2 PENCIL only.
- Fill in only ONE BUBBLE for each part of a question.
- · Be sure to read each question carefully.

EXAMPLE QUESTION







SERIAL #

1. Are you a: Boy Girl O O 2. What grade are you in? 4 5 6 O O O	9. During the current school year, do you qualify for a free or reduced price school lunch? Yes No Don't Know O O O 10. Do one or both of your parents usually attend school-sponsored open houses or PTA meetings? Yes No O O
3. Do you live with both of your parents? Yes No	Below is a list of things some people sniff to get high. These are called inhalants. 11. Have you EVER sniffed any of the following
4. How old are you? 8 or younger 9 10 11 12 ofder O O O O O O 5. Are you: African Mexican Asian Native American Other O O O O O 6. What grades do you usually get? (ONLY CHOOSE ONE) Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's O O O O	inhalants to get high? (DARKEN ONE BUBBLE FOR EACH DRUG) Never Used Used a. Gasoline to get high b. Paint thinner to get high c. Glue to get high d. Whiteout or correction fluid to get high e. Spray paint to get high f. Other inhalants to get high The following questions ask about drugs people use to get high. 12. How many times have you used any of these drugs? (DARKEN ONE BUBBLE FOR EACH DRUG)
7. Have you lived in this town for more than 3 years? Yes No Don't Know O O 8. Did either of your parents graduate from college? Yes No Don't Know O O O	a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron? h. Inhalants (whiteout, glue, gas, etc.)? i. Marijuana (pot)?

13. SINCE SCHOOL BEGAN IN THE FALL, how many times have you used:	15. About how many of your close friends your age use:
(DARKEN ONE BUBBLE FOR EACH DRUG)	(900-99)
Times used since school	(DARKEN ONE BUBBLE FOR EACH DRUG)
began in the Fall.	
	.::/ / /
± 8	Never heard of it None Some
Ore Ses A it	
fin lise let	Never None Some
Never heard of it Not used it 1-2 times 3-10 times	a. Cigarettes?
a. Cigarettes?	b. Snuff or Chewing
b. Snuff or Chewing	Tobacco?
Tobacco?	Tobacco?
c. Beer?	d. Wine Coolers?
Tobacco?	
e. Wine? OOOO	f. Liquor (whiskey, vodka, tequila, etc.)?
	tequila, etc.)?
vodka, tequila, etc.)?	h. Inhalants (whiteout, glue,
h. Inhalants (whiteout,	gas, etc.)? O O O
glue, gas, etc.)?	i. Marijuana (pot)?
i. Marijuana (pot)?	
14. How old were you when you first used:	16. Has anyone ever tried to give you:
(DARKEN ONE BUBBLE FOR EACH DRUG)	(DARKEN ONE BUBBLE FOR EACH DRUG)
IDAINEN ONE BOBBLE I ON EACH BIDGY	(DATRET ONE BOBBLE FOR EACH BROOK
d of h	<u> </u>
Never heard of Never heard of 7 or younger 8 years 10 years 11 years 12 years or of 12 years or of 12 years or of 13 years or of 14 years or of 14 years or of 15 years or of 15 years or of 16 years or of 17 years or of 18 years or	9/ /
Never heard Never used: 7 or younges 8 years 10 years 11 years 12 years or	hee
/ ½ / ½ / ½ / ½ / ½ / ½ / ½ / ½ / ½ / ½	~ 1 1 1
0101012121-1-1-1	20 20 20 20 20 20 20 20 20 20 20 20 20 2
	Never heard of it
a. Cigarettes?	a. Cigarettes?
a. Cigarettes? b. Snuff or Chewing	a. Cigarettes? O O b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco?	a. Cigarettes? O O b. Snuff or Chewing Tobacco? O O
a. Cigarettes? b. Snuff or Chewing	a. Cigarettes? O O b. Snuff or Chewing Tobacco? O O
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer?	a. Cigarettes? O O O b. Snuff or Chewing Tobacco? O O O c. Beer? O O
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka,	a. Cigarettes? O O O b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)?	a. Cigarettes? O O O b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron?	a. Cigarettes? b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron? h. Inhalants (whiteout,	a. Cigarettes? b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron?	a. Cigarettes? b. Snuff or Chewing
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron? h. Inhalants (whiteout, glue, gas, etc.)?	a. Cigarettes? b. Snuff or Chewing

PLEASE GO ON TO THE NEXT PAGE

17. When you drink alcohol (beer, wine coolers, wine or liquor), where do you get it? (DARKEN ONE BUBBLE FOR EACH LINE) Don't Drink Yes No a. Get it from home OOO b. Get it from friends OOO c. Get it from somewhere else	20. During the past year (12 months), how many times have you had two or more glasses or bottles of these drinks in a row? (DARKEN ONE BUBBLE FOR EACH DRUG) 1 2 3 or Never time times more times a. Beer O O O O b. Wine Coolers O O O c. Wine O O O d. Liquor O O O
18. How dangerous do you think it is for kids your age to use: (DARKEN ONE BUBBLE FOR EACH DRUG)	21. Since school began in the Fall, have you skipped school when your parents didn't know?
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Shimeron? h. Inhalants (whiteout, glue, gas, etc.)? i. Marijuana (pot)?	22. Since school began in the Fall, have you been sent to anyone like the principal because you did something against the rules? O Yes O No 23 How do your parents feel about kids your age drinking beer? They don't like it. They don't care. They think it's O.K. I don't know.
19. SINCE SCHOOL BEGAN IN THE FALL, have you learned about drugs or alcohol from: (DARKEN ONE BUBBLE FOR EACH LINE) Yes No a. Your teacher? b. A visitor to your class? c. An assembly program? d. A school counselor? e. Someone else at school?	24. How do your parents feel about kids your age using marijuana? They don't like it. They don't care. They think it's O.K. I don't know.
	for being part of this important project.

PLEASE DO NOT WRITE IN THIS AREA

SERIAL#

Mark Reflex® forms by Pearson NCS MM96595-8 654321 ED05 Printed in U.S.A.

TEXAS SCHOOL SURVEY OF SUBSTANCE USE

SECONDARY SURVEY INSTRUMENT 2006 VERSION

INTRODUCTION

Before answering the survey questions in this booklet, please read the following:

- I am aware that the survey questions I am answering are designed to measure drug and alcohol use in my school district and the state of Texas, that I will be asked questions about student attitudes and behaviors that relate to substance abuse issues, and that over 300,000 students all over Texas will be taking this survey;
- I know that I do not have to take this survey and that I can skip any question that I do not choose to answer for any reason without any consequences;
- I know that there is no way for anyone to find out which survey that I answered, that the numbers on the questionnaire books are used only to identify which pages go together after the booklets are cut and cannot be used to tell who answered the questions; and
- > I am voluntarily participating in the Texas School Survey of Drug and Alcohol Use.

If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol or drugs, counselors from the Worker's Assistant Program of Texas are available by calling this toll free number: 1-877-9-NO-DRUG (1-877-966-3784).

This research study has been reviewed and approved by the Institutional Review Board — Human Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted through Ms. Angelia Raines, Director of Research Compliance, Office of the Vice President for Research at (979) 458-4067 or email at araines@vprmail.tamu.edu.

DIRECTIONS

- · DO NOT write your name anywhere on this booklet.
- · Use a NUMBER 2 PENCIL only.
- Fill in only ONE BUBBLE for each part of a question.
- · Be sure to read each question carefully.

EXAMPLE QUESTION

IN	THE PAST WEEK, on how m	any DAYS hav	e you us	ed:		
		Never heard of it	Not used It	1-2 days	3-4 days	5-7 days
a.	Cigarettes	0	0	0	0	
b.	Smokeless Tobacco	0	0		0	0
c.	Beer	0		0	0	0





OCCOCCOCCOCCOCCOCCOCCOCCOCCOCC

SERIAL #

1. Are you:	7. How long have you attended school in this District?
Male Female	1 2-3 4 or year or less years more years
2. What grade are you in? 7 8 9 10 11 12 O O O O	8. Do you have a job? Yes No O O
3. Do you live with both of your parents? Yes No O O	9. Do you get an allowance? Yes No
4. How old are you? 11 or younger 12 13 14 15 16 17 18 19	10. Did either of your parents graduate from college? Yes Don't Know
5. Are you: African Mexican Asian Native White American American American Other OOOOOOO	11. During the current school year, do you qualify for a free or reduced price school lunch? O Yes
6. On average what grades do you get? (DARKEN ONE BUBBLE ONLY) Mostly Mostly Mostly Mostly A's B's C's D's F's O O O	O Don't Know

12.	Do you reg	ularly particip	ate in any of the following	extra-curric	ular SCHOOL	activities?	(DARKEN C	NE BUBBLE FOR	EACH LINE: a-m.)				
	Ye	s No											
	a. C		School Athletics										
	b. C		School Band/Orchestra										
	c. O School Choir												
		O Drama, Speech, or Art Clubs											
	e. C		O Drill Team or Cheerleading Squad										
	f. C		Student Government										
	g. C		Student Newspaper or '	tudent Newspaper or Yearbook									
	h. C		Academic Clubs or Soci	eties (math,	, science, for	eign lang	uage, etc.)						
	i. C		Academic Competition	Groups (UIL	., science bo	wl, etc.)							
	j. C		Service Clubs at School	(Key Club,	Peer Mediate	ors, PALS)							
	k. C		Other School Clubs or S	tudent Gro	ups								
	I. C		Athletic Teams OUTSID	E of School									
	m. C		Other Clubs or Groups (OUTSIDE of	School								
12	How safe	do vou feel w	hen you are:										
10.			FOR EACH LINE: a-c.)										
	DAIRCEN	ONE BODDEL	TOTT EACH EINE. a c.,	Very	Somewhat	Not V	ery N	Not Safe	Don't				
				Safe	Safe	Saf	e	At All	Know				
	a. in your	home		0	0	O		0	0				
	b. out in	our neighbo	rhood	0	0	10		0	0				
	c. at scho	ol		0	0 1	_0)	0	0				
						1							
14.	a. Feel clo	ONE BUBBLE	ENDS would you say: EFOR EACH LINE: a-e.) Parents? Papons like a knife or gu good grades?	None O	A Few		Some O O	Most O O	All O O				
			are interested in becom				O		0				
		member?	are interested in become				0						
	The second secon		op out of school?	00	0		ő	0	00				
15.	Do one or	both of your	parents usually attend s	chool-spons	sored open h	nouses							
	or PTA me		The second secon				01	res OI	No				
									-				
	Below is a	list of things	some people sniff to ge	t high. They	are called II	NHALANT	S.						
16.			ave you sniffed, huffed, o	or inhaled th	ne following	INHALAN	TS for "kid	cks" or to get	"high":				
	DARKEN	JNE BUBBLE	FOR EACH LINE: a-k.)					Used at					
						ver ard	Used at Least once	Least once Since Schoo	Used at Least once				
						lever	In the	Began in	In Your				
							Past Month		Lifetime				
	a. Liquid	or spray pair	nt to get high?)	0	0	0				
	b. Whiteo	ut, correctio	n fluid to get high?)	0	0	0				
	c. Gasolii	ne to get hig	h?				0	Ö	0				
		o get high?					0	Ö	0000000000				
			om, Rush, Bolt, Climax to	get high?)	0	Ö	Ō				
	f. Glue to	get high?		-			0	0	O				
	g. Paint o	r lacquer thi	nner, toluene, other solve	ents to get l	high? (0	O	0				
			hing gas, whippets) to ge)	0	0	0				
		booster to					0	0	0				
	j. Other s	prays (non-s	stick pan coatings, hair sp	oray, etc.) to	get high? (0	Ŏ	0				
		nhalants to			(0	0	0				

A-8

SERIAL#		OOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOO						
17. How recently, if ever, have you used the TOBACCO and ALCOHOL products ever (DARKEN ONE BUBBLE FOR EACH LIN	en one time:	(DARKEN ONE BUBBLE FOR EACH LINE: a-h.)						
/	a. b. c. d. e. f. g.	Cigarettes? Smokeless Tobacco? Beer? Wine Coolers? Wine? Liquor? Inhalants (whiteout, glue, gas, etc.)?	0 000000 None heard of					
18. How recently, if ever, have you used the DRUGS even one time: (DARKEN ONE B LINE: a-k.) LINE: a-k.)	UBBLE FOR EACH 21. I	DARKEN ONE BUBBLE FO	/////	/				
a. Marijuana? b. Cocaine (not crack)? O c. Crack? d. Shimeron? e. Hallucinogens (LSD, PCP, Mushrooms, etc.)? O f. Uppers? g. Downers? h. Rohypnol (roches, roofies, etc.)? i. Steroids? j. Ecstasy? k. Heroin?	a. b. c. d. e. f. g. h. i. j.	Cigarettes? Osmokeless Tobacco? Osmokeless Tobacco. Osmokeless Tob	00000 00000	Oash				
a. Tobacco? b. Alcohol? c. Inhalants (whiteout, glue, gas, etc.)? d. Marijuana? e. Cocaine (not crack)?	es, if any, have EACH LINE: a-f.) n. o.	Hallucinogens (LSD, PCP, Mushrooms, etc.)? Uppers? Downers? Rohypnol (roches, roofies, etc.)? Steroids? Ecstasy? Heroin?						

	SERIAL#	VOOOOOOOOO								
22.	How old were you wh (DARKEN ONE BUBBI					r.)			1	24. During the past 30 days, how many cigarettes have you smoked?
a.	Cigarettes?		O Never us	O o rounger		200		00 1% 00 1%		
b. c. d.	Smokeless Tobacco? Beer? Wine Coolers? Wine?		0000							a. Cidarettes ber day hore day
e. f.	Liquor?		0	300				30		a. Cigarettes? O O O O O
1000	Inhalants (whiteout,								1	
	glue, gas, etc.)?		00					OC		25. When you drink alcoholic beverages, how many drinks do you usually have AT ONE TIME, on
h.	Marijuana?		0					OC		average:
i.	Cocaine (not crack)?		OK					SIO		(DARKEN ONE BUBBLE FOR EACH LINE: a-d.)
j.	Crack?		O							
k.	Shimeron? Hallucinogens (LSD,		00							Never drink this beverage 12 or move drinks 5 8 drinks 2 drinks 1 drinks 1 drinks 1 drinks 1 drinks
- 54	PCP, Mushrooms, etc.)	?	00					00	-	Never drink This beverage 12 or more drinks 9 - 1 drinks 2 drinks 2 drinks 1 drink Less than 1 drink
m.	Uppers?		O					SO	(Never drink this beverage 12 or more art 3 4 drinks 3 4 drinks 1 drink Less than 1 dr
n.	Downers?		0					OC		Never drii This bever 12 or mon 3 - 4 driinks 1 driinks Less than
0.	Rohypnol (roches,			1		1	1//			1 1 2 3 3 3 3 3 3 3 4 5 5 5 5 5 5 5 5 5 5 5 5
	roofies, etc.)?		-OK	51616	SIÓIG	5 0		50/	V	\ \ a. \ Beer \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \
p.	Steroids?	1	2	3					1	b. Wine Coolers O OOOOO
q.	Ecstasy? Heroin?	1	8	3				XIXI		c: Wine
r.										
23.	How often do you no				1	11	7			26. IN THE PAST 30 DAYS, on how many days
	(DARKEN ONE BUBBI	EFC	H EA				/	1 9	b	have you had 5 or more drinks of alcohol AT ONE TIME?
	Cigarettes?	O Never us	O Every da.	Several time	Several time	About once	About onco	O Less than once		\$ / / / / / /
a. b.	Smokeless Tobacco?	o	0	0	0	0	0	ŏ		day,
c.	Beer?	ŏ	ŏ	00	ŏ	ŏ	ŏ	ŏ		Never had 5 or m thinks at one time None 7 day 2 days 3 to 5 days 6 to 9 days 10 or more days
d.	Wine Coolers?	Ö	Ö	0	O	Ö	0	Ö		Never h frinks an None T day 2 days 3 to 5 di 6 to 9 d
e.	Wine?	0	0	00	0	0	00	0		10 6 t 3 t 20 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
f.	Liquor?	0	Ŏ	0	0	O	0	0		a. Alcohol O O O O O
g.	Inhalants (whiteout,	_								
b	glue, gas, etc.)? Marijuana?	00	0000	00000	00000	00000	00000	000		
h. i.	Cocaine (not crack)?		0	0	0	0	0	0		27. How often do you get alcoholic beverages from
j.	Crack?	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ		the following sources?
k.	Shimeron?	Ŏ	00	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ		(DARKEN ONE BUBBLE FOR EACH LINE: a-e.)
I.	Hallucinogens (LSD,									<u> </u>
	PCP, Mushrooms, etc.)	?0	0	0	0	0	0	0		
50000	Uppers?	0	0	00	0	0	0	0		a. At home
n.	Downers?	0	0	0	0	0	0	0		2 At home
0.	Rohypnol (roches, roofies, etc.)?	0	0	0	0	0	0	0		a. At home OOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOO
p.	Steroids?	Ö	ő	0	0	00	ŏ	ŏ		c. From a store
q.	Ecstasy?	O	ŏ	O	ŏ	Ö	ŏ	ŏ		c. From a store
r.	Heroin?	0	O	O	O	0	O	0		e. Other source

28.	How often do you use marijuana in the following ways? (DARKEN ONE BUBBLE FOR EACH LINE: a-e.) a. Joints b. Blunts c. Bongs or water pipes d. Pipes e. Other ways		000	8 8 8	OOOOO BON'S KROW
29.	How recently, if ever, have you taken the following over-the-counter or prescription-type drugs to get "high": (DARKEN ONE BUBBLE FOR EACH LINE: a-b.) a. DXM, Triple C's, Skittles, or Coricidin to get high? b. Codeine cough syrup (Lean, Nods, AC/DC) to get high?	rd Least or over In the lit Past Mo	nt L	Used at east once nce School Began in the Fall	Used at Least once In Your Lifetime
	How dangerous do you think it is for kids your age to use: (DARKEN ONE BUBBLE FOR EACH LINE: a-i.) a. Tobacco? b. Alcohol? c. Inhalants? d. Marijuana? e. Cocaine (not crack)? f. Crack? g. Ecstasy? h. Steroids? i. Heroin?		0000000		/9 /
31.	since school began in the fall, on how many DAYS have you (DARKEN ONE BUBBLE FOR EACH LINE: a-g.) a. missed a whole day of school because you "skipped" or "cut"? b. missed a whole day of school because you were ill? c. missed a whole day of school for some other reason? d. been sent by a teacher to someone like the Principal, Dean, or Guidal Counselor because of your conduct or attitude? e. had someone from your home be called to school because of your conduct or attitude? f. gotten into trouble with your teachers because of your drinking? g. gotten into trouble with your teachers because of your drug use?	None O O O O O O O	1-3 days O O O	days O O O O	10+ days O O O O
32.	DURING THE PAST 12 MONTHS, how many times (if any) have you (DARKEN ONE BUBBLE FOR EACH LINE: a-h.) a. gotten into difficulties of any kind with your friends because of your drinking? b. driven a car when you've had a good bit to drink? c. been criticized by someone you were dating because of drinking? d. gotten into trouble with the police because of drinking?	None O O O	1-3 times	4-9 times	10+ times
	e. gotten into difficulties of any kind with your friends because of your drug use? f. driven a car when you've felt high from drugs? g. been criticized by someone you were dating because of drug use? h. gotten into trouble with the police because of drug use?	0000	0000	0000	0000

33.	SINCE SCHOOL BEGAN IN THE FALL, on how many DAYS (if any) have you attended at least one class while "high," "drunk" or "stoned" on										
	(DARKEN ONE BUBBLE FOR EACH LINE: a-d.)			None days		days	4-9 days		10+ days		
	a. beer, wine coolers, wine, or hardb. marijuana?	liquor?				0	0				
	c. inhalants?				5	0	Ö		ŏ		
	d. some other drug(s)?				ŏ	ŏ	ŏ		ŏ		
34.	Alcohol or other drugs are sometimes used at teenage parties. Thinking of the parties you attended this school year										
	(DARKEN ONE BUBBLE FOR EACH LI	INE: a-b.) Never	Seldom	Half the time	Most of the time	Always	Don't Know	Didn't att			
	a. how often was alcohol used?	0	0	0	0	0	0	0			
	b. how often were marijuana	0			0	0		0			
	and/or other drugs used?	0	0	0	0	0	0	0			
35.	SINCE SCHOOL BEGAN IN THE FALL, have you sought help, other than from family or friends, for problems in any way connected with your use (if any) of alcohol, arrijuana, or other drugs?								-		
36	If you had a drug or alcohol problem	and needed	- (37. SIN	CE SCHOOL	L BEGAN IN	THE FALL	have vo			
00.	help, who would you go to?		11	1.1		ormation on					
	(DARKEN ONE BUBBLE FOR EACHLE		es No	the	following S	CHOOL sou	ırces?		-		
	a. A counselor or program in school? O O DARKEN ONE BUBBLE FOR EACH LINE: a-h							and the second second			
	 Another adult in school (such as or teacher)? 		0 /0	1	A school he	alth class?		Yes	No O		
	c. A counselor or program outside		5 0	- Contraction		ly program	at school?	_	ŏ		
	d. Your parents?	11/8				idance cou		O	0		
	e. A medical doctor?	1	5 0		A science c			0	0000		
	f. Your friends? g. Another adult (such as relative,			DOLD INC	A social stu	idy class? up or club mee	ting of color	017 ()	\approx		
	clergyman, or other family friend	1)? (0			school gues		0	ŏ		
	h. I wouldn't go to anyone.					urce at scho		ŏ	ŏ.		
\vdash											
38.	SINCE SCHOOL BEGAN IN THE FALL	The second secon	otten any in	formation	on drugs o	r alcohol fro			_		
	(DARKEN ONE BUBBLE FOR EACH L	INE: a-d.)					Yes		No -		
	a. A website or online chat group?	mark lada 2					00	(2		
	 Books, magazines, posters, or pa TV, radio, or other audio/video? 	mpnietsr					0	(
	d. Your parents?						ŏ	(ŏ •		
39.	How do your parents feel about kids	your age	(DARKEN	ONE BUBE	BLE FOR EA	CH LINE: a-c	c.)		-		
	Strongly	Mildly		er Approve	Mildly		trongly		on't		
	a. smoking cigarettes?	Disappro	ve nor E	Disapprove	Approv	re A	O		O		
	b. drinking beer?	ŏ		ŏ	ŏ		ŏ		ŏ		
	c. using marijuana?	Ö		ŏ	Ö		Ö	(Ŏ -		
40	In the past school was because to	n tooked for	duna maa bu		ot onbession		O V		O No		
40.	In the past school year, have you bee	n tested for	urug use by	someone	at school?		O Yes) No		
41.	IN THE PAST YEAR, have you purcha	sed a Texas	Lottery tick	et?			O Yes	(○ No		

SUPPLEMENTAL SURVEY QUESTIONS S-4 (A) S-7 (A) S-8 (A) S-9 (A) S-1 (A) S-2 (A) S-3 (A) S-10 (A) 00000000000 B 00000000000 **B** @@@@@@@ @@@@@@ 0 0000000000 0 (E)(C)(E) 0 0 0 0 0 0 S-13 (A) S-14 (A) (A) (A) S-11 (A) **B** @@@@@@@ @©@@@@@ @@@@@@@ @@@@@@ **B** 000000000 0 0 0 S-28 (A) S-22 (A) S-23 (A) (A) A **EODODO** 00000 A OOOOOOOOOOS-26 @@@@@@ @@@ 00000000000 00000000000 ₿ (B) 90000 0 0 900 0 @@@ 000 0000 **©** (E) 0 O 0 0

THANK YOU FOR BEING PART OF THIS IMPORTANT PROJECT

PLEASE DO NOT WRITE IN THIS AREA

SERIAL #

Mark Reflex® forms by Pearson NCS MM96596-9 654321

ED05

Printed in U.S.A.

APPENDIX B Participating Districts

2006 Participating Districts

DISTRICT	NO.	STRATA	
PLAINVIEW	095905		12
FRENSHIP	152907		13
SHALLOWATER	152909		13
SHAMROCK	242902		13
NORTHSIDE	015915		23
MIDWAY	161903		23
ARLINGTON	220901		31
FORT WORTH	220905		31
PLANO	043910		32
GARLAND	057909		32
GRAND PRAIRIE	057910		32
MAYPEARL	070915		32
SHERMAN	091906		32
GRANBURY	111901		32
CORSICANA	175903		32
ALLEN	043901		33
DENISON	091903		33
GRAPEVINE COLLEYVILLE	220906		33
MANSFIELD	220908		33
CROWLEY	220912		33
BRIDGEPORT	249903		33
AUBREY	061907		34
ALEDO	184907		34
CHISUM	139905		42
EDGEWOOD	015905		43
HOOKS	019902		43
KILGORE	092902		43
LONGVIEW	092903		43
EUSTACE	107905		43
NEW DIANA	230906		43
BEAUMONT	123910		51
NACOGDOCHES	174904		52
CYPRESS-FAIRBANKS	101907		61
HOUSTON	101912		61
CLEAR CREEK	084910		62
SPRING BRANCH	101920		62
WACO	161914		62
CONROE	170902		62
COLUMBIA - BRAZORIA	020907		63
EAST CHAMBERS	036903		63

DEER PARK	101908	63
NORTH FOREST	101909	63
LA PORTE	101916	64
LIBERTY	146906	64
ROUND ROCK	246909	71
GEORGETOWN	246904	72
LEANDER	246913	72
COLLEGE STATION	021901	73
ROSEBUD LOTT	073905	73
MILANO	166903	73
SAN ANTONIO	015907	81
BOERNE	130901	82
SAN FELIPE DEL RIO	233901	82
JUDSON	015916	83
MEYERSVILLE	062906	83
NURSERY	235904	83
FLORESVILLE	247901	83
SEGUIN	094901	84
SAN ANGELO	226903	92
ALPINE	022901	102
CLINT	071901	102
SAN ELIZARIO	071904	102
YSLETA	071905	102
SOCORRO	071909	102
BROWNSVILLE	031901	112
LOS FRESNOS	031906	112
RIO HONDO	031911	112
SOUTH TEXAS	031916	112
EDCOUCH-ELSA	108903	112
McALLEN	108906	112
MERCEDES	108907	112
PHARR-SAN JUAN-ALAMO	108909	112
WESLACO	108913	112
LA VILLA	108914	112
MONTE ALTO	108915	112
VALLEY VIEW	108916	112
RIO GRANDE CITY	214901	112
ROMA	214903	112
LYFORD	245902	112
RAYMONDVILLE	245903	112
PETTUS	013903	113

APPENDIX C

Classroom Selection Guidelines

Texas School Survey of Substance Use

Teacher Lists for Random Sample Selection

For elementary

- List <u>all</u> the teachers' names that teach in grades 4, 5 and 6;
- List the grade level that each teaches; and
- List the number of students per class.

For secondary

- Choose *either* a class period in which <u>all</u> students are present *or* a subject all students must be enrolled;
- List <u>all</u> teachers for that class period or that subject; and
- List the number of students *per grade level* in each of those classes.

Note:

If a subject has been chosen rather than a particular class period, and where a single teacher teaches multiple classes, a classroom identifier of some sort (e.g., the period) should be included in the column under the heading of "period".

PPRI will randomly pick entire classrooms until we reach the target numbers per grade level for the sample. Then, we'll send you the survey materials along with a Master List of those classes randomly selected to be surveyed.

Helpful Hint:

Electronic submissions (i.e., as an attachment to an email) are <u>encouraged</u> for speedy turnaround, as the random sample selection is a computer program.

Remember:

Sixth graders are considered elementary students for the purposes of this survey.

APPENDIX E

School Survey Administration Materials

District Coordinator's Responsibilities

Campus Coordinators' Responsibilities

Teachers' Guide

Survey Administration Procedure Survey Administration Guide

Texas School Survey of Drug and Alcohol Abuse

District Coordinator's Responsibilities

1. Distribution of the Survey Administration Guide

The *Survey Administration Guide* enclosed in this mailing list is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy.

The guides should be sent to the campuses in the time to be distributed one week prior to the scheduled survey administration date. This will allow time for everyone involved to read through the information and resolve any questions.

2. Distribution of Survey Instruments and Envelopes

Enclosed is a set of envelopes. There should be approximately one envelope per classroom, according to the number of classrooms indicated on the survey enrollment form. The envelopes will be used to package the blank survey forms sent to each class and seal the completed instruments after survey administration (see Campus Coordinators' Responsibilities).

At least one week prior to the scheduled survey date, District Coordinators should forward the appropriate number of elementary and secondary survey instruments and envelopes to each campus. Each campus will need as many envelopes as there are classes participating. In case of mass administration, approximately one envelope per 24 students should be used.

3. Packaging the Completed Surveys for Delivery to PPRI

Completed survey instruments will be sealed in envelopes and collected from each classroom. As the District Survey Coordinator, you should clearly designate to each Campus Coordinator where the surveys should be sent from there.

All completed surveys should be collected at a central location. Completed surveys should remain in the sealed envelopes, and the envelopes from each campus should be grouped together and **CLEARLY LABELED** with the campus name. **BLANK SURVEYS SHOULD BE** returned to PPRI along with the classroom envelopes. The survey materials should be boxed, and the boxes should be labeled sequentially (i.e., "one of three, two of three," etc.). Materials should then be returned to PPRI in a single mailing. Ship to:

Texas School Survey
PPRI / Texas A&M University
317 Dulie Bell Bldg. College Station, TX 77843-4476

Texas School Survey of Drug and Alcohol Abuse

Campus Coordinators' Responsibilities

1. Distribution of the Guide to Survey Administration

Each Campus Coordinator will receive copies of the Survey Administration Guide and Procedure from the District Coordinator. The Guide should be distributed to all survey administrators approximately one week prior to the scheduled administration date so that everyone will have adequate time to read through the information and resolve any questions.

2. Distribution of Survey Instruments to Classes

The District Coordinator will send to each participating campus the appropriate number of survey instruments and enough envelopes to provide one to each class being surveyed. When the materials arrive, Campus Coordinators should follow these steps:

- a) Count out a set of surveys for each participating class
- b) Place each set of surveys in an envelope
- c) Label each envelope to indicate which class it is along with **the campus name** and how many survey instruments are enclosed. DO NOT seal the envelopes. DO NOT remove the classroom information form stapled to each envelope.

On the day the survey is to be given, envelopes filled with the correct number of survey instruments should be distributed to each survey administrator. The individuals administering the survey should have reviewed the *Survey Administration Guide and Procedure* (described above) prior to this point and should already be aware of their responsibilities.

3. Collecting Completed Survey Instruments

After the survey has been administered in each class, teachers will be instructed to seal the completed survey forms in the envelopes along with the completed classroom identification form. Unused blank forms should be kept together **OUTSIDE THE ENVELOPE**. It is recommended that teachers not leave the room with the sealed envelopes. Rather, someone who did not participate in the actual survey administration should be sent around to collect the envelopes.

Once envelopes have been collected from every classroom, they should be forwarded to the location designated by the District Survey Coordinator where they will be collected and returned to PPRI in a single mailing.

Texas School Survey of Drug and Alcohol Use

Survey Administration Guide

Your school district has elected to participate in the Texas School Survey of Drug and Alcohol Use. The survey is sponsored and designed by the Texas Commission on Alcohol and Drug Abuse (TCADA) and is implemented by the Public Policy Research Institute (PPRI) at Texas A&M University. The district needs your assistance in administering the survey. PPRI has prepared this Survey Administration Guide to provide information and instructions for conducting the survey project.

Your classroom envelope should contain blanks surveys and an attached Classroom Identification Form to be filled out during survey administration.

Please read all instructions carefully before administering the survey. It is important that outlined procedures are followed so that administration of the survey is uniform throughout the district.

Your attitude towards this survey is extremely important. If students sense the importance of this information, they will take the survey more seriously. Please stress to the students that their responses are entirely anonymous.

In addition, your physical position in the classroom may influence the way students answer questions. While students are taking the survey, do not walk around the room. Please sit or stand in a part of the room away from students.

Try to keep students as quiet as possible throughout survey administration. If students are allowed to talk about the contents of the survey or their own experiences, peer influence could cause students to either inflate or under-report their own drug and alcohol habits.

The survey should take approximately 45 minutes to administer.

To maintain student confidentiality, do not view completed survey instruments.

Students must use a #2 pencil. Be prepared to provide pencils to students who may not have one. Surveys completed in pen or marker will not be processed.

Survey Administration Procedure

1. Before handing out surveys, tell students:

This survey is for the Public Policy Research Institute at Texas A&M University. Survey results will be used for evaluating and planning substance abuse programs for our school district. No student is required to participate in this survey. This survey will ask questions about your experiences with drugs and alcohol. Do not turn the pages or begin answering questions until we have gone over the survey instructions.

- 2. Make sure all students have a #2 pencil to complete the survey.
- 3. Hand out surveys and tell students:

I cannot answer questions about specific substances or tell you what questions mean. If you are not familiar with a substance listed on the survey, mark the category "Never heard of it."

- 4. Then go over the instructions on the attached survey form, using either the elementary or secondary form. Be sure to read the Introduction, Directions, and Example Question.
- 5. Emphasize the following points to the students:
 - **This survey is completely voluntary.
 - **No one will know how individual students answer questions.
 - **If you do not feel comfortable answering a question or feel you cannot answer a question honestly, leave it blank.
 - **A drink means a can or bottle of beer or wine cooler, a 4 ounce glass of wine, a shot of liquor, or a mixed drink.
- 6. If students refuse to take the survey, have them work quietly at their seat or send them to an area designated by your campus coordinator.
- 7. Collect all completed surveys and place in envelope. **DO NOT PUT UNUSED SURVEYS IN THE ENVELOPE**. Your district will not be reimbursed for blank surveys left in the envelopes. Do not view completed surveys instruments.
- 8. Fill out the attached Classroom Identification Form and place inside the envelope. Seal the envelope.
- 9. Return classroom envelope and unused surveys to your campus coordinator.

APPENDIX F

Computation of Sample Weights

Weighting the Texas School Survey of Substance Abuse

District Level Weights

Within the selected districts, students were sampled via a simple random sample of classrooms (i.e., a sample in which all classrooms in the district had an equal chance of getting in) picked directly from the district. Every student in the sampled classrooms was asked to participate in the survey.

A weight w_{ilk} is assigned to each sampled classroom k, in grade l, in district i:

$$w_{ilk} = \frac{N_{il}}{n_{il}} \tag{1}$$

where.

 N_{il} = total no. of classrooms in grade l, in district i

 n_{il} = no. of classrooms sampled in grade l, in district i

Since every student in the sampled classrooms was included in the sample, each student's probability of selection is equal to his/her classroom's probability of selection. Thus, each sampled student in these districts is assigned his/her classroom weight. Therefore the weight w_{ilr} of the r^{th} sampled student in classroom k is,

$$W_{ilr} = W_{ilk} \tag{2}$$

<u>Remark</u>: In (1), if the true N_{il} is not known for a particular district, estimate it with,

$$\hat{N}_{il} = \frac{M_{il}}{\hat{c}_{il}} \tag{3}$$

where, $\hat{c}_{il} = m_{il} / n_{il}$ and

 M_{il} = total no. of students in grade l, in district i

 m_{il} = no. of students sampled in grade l, in district i

Notice that in this case, the weights in (1) reduce to, $w_{ilk} = M_{il} / m_{il}$.

District Level Estimates

The total no. of students in grade l, in district i, giving a positive response to a question is estimated as

$$\hat{Y}_{il} = \sum_{r=1}^{m_{il}} w_{ilr} y_{ilr} \tag{4}$$

where

$$y_{ilr} = \begin{cases} 1 & \text{if the } r^{\text{th}} \text{ student's response is positive} \\ 0 & \text{otherwise} \end{cases},$$

and w_{ilr} is as defined in (2).

The proportion of students in grade l, in district i, giving a positive response to a question is estimated as,

$$\hat{\overline{Y}}_{il} = \frac{\hat{Y}_{il}}{M_{il}}$$

Stratum Level Weights

The districts were selected from strata that were formed on the basis of two factors, namely, how urban the districts were and the levels of enrollment. That is, once the districts were grouped by the degree to which they were urban, strata were formed by collecting together districts of similar size within these urban classes. Next, a simple random sample of districts was chosen from each stratum. A few large districts were selected to the sample with probability one. These districts were grouped into two strata based on the relative sizes of the districts.

Stratum level weights are assigned to the sampled students as follows. The weight w_{min}^{s} of the r^{th} sampled student, from grade l, in district i in stratum h is,

$$w_{hilr}^{s} = \frac{1}{p_{hi}} w_{ilr} \tag{5}$$

where

$$p_{hi} = \begin{cases} 1 & \text{if district } i \text{ was sampled with probability 1} \\ \phi_{hi} & \text{otherwise} \end{cases}, \tag{6}$$

and within grade l, for each district i in stratum h, w_{ilr} is as defined in (4), and $\phi_{hi} = n_h / N_h$ with

 n_h = no. of districts sampled from stratum h

 N_h = total number of districts in stratum h

Notice that within stratum h, ϕ_{hi} does not depend on district i.

State Level estimates

The total no. of students in grade l, giving a positive response to a question is estimated as

$$\hat{Y}_l = \sum_{h=1}^H \hat{Y}_{hl} \tag{7}$$

with

$$\hat{Y}_{hl} = \sum_{i=1}^{n_h} \sum_{r=1}^{m_{il}} w_{hilr}^s y_{ilr}$$
 (8)

where, within grade l, for each district i in stratum h, y_{ilr} , and m_{il} are as defined above.

The proportion of students in grade l, in the state, giving a positive response to a question is estimated as,

$$\hat{\bar{Y}}_{l} = \frac{1}{M_{l}} \sum_{h=1}^{H} \hat{Y}_{hl} \tag{9}$$

where,

 M_l =total number of students in grade l in state.

APPENDIX G

Standard Error Computation

Standard Error Computation for the Texas School Survey of Substance Use

The standard errors of the percentages were estimated via the SUDAAN software package. The variance estimation techniques in SUDAAN are specifically designed to address the complexity of the sample design in finite population sampling. The variance estimators generally available for these complex designs do not have closed form solutions and, therefore, require iterative approaches in computations that would yield solutions as close to the true value as possible. SUDAAN software package is available for this purpose and is gaining wide popularity in survey sampling applications.

In applying SUDAAN to the current survey which consisted of a multi-stage stratified sampling design, the input SAS data set requires the following re-coding of the stratification variable for strata 21 and 22 (due to certainty stratum 21 having only one district responding to survey):

```
if strata=22 then strata=21;
(Here: strata= stratum identifier) Next, the data set needs to be sorted as follows:
```

```
proc sort data=___;
by strata distid q2 class;
run;
```

Here, distid= district identifier, q2= grade identifier, class= classroom identifier.

```
The SUDAAN code needs to incorporate the following: proc crosstab data=" " filetype=sas design=WOR; nest strata distid q2 class;
```

The specified design WOR uses Taylor series variance estimation methods¹ in computing standard errors of percentages in the multi-stage stratified sampling design of this survey.

_

¹ See SUDAAN User's Manual (Volume 1), page 3-4.

APPENDIX H

Standard Error and Confidence Interval Tables

Elementary Grades – 2006

Standard Error and Confidence Intervals

All Students-Overall and by Grade

Past Year Use

Lifetime Use

Male-Overall and by Grade

Past Year Use

Lifetime Use

Female-Overall and by Grade

Past Year Use

Lifetime Use

Whites-Overall and by Grade

Past Year Use

Lifetime Use

African American-Overall and by Grade

Past Year Use

Lifetime Use

Mexican American-Overall and by Grade

Past Year Use

Lifetime Use

Border-Overall and by Grade

Past Year Use

Lifetime Use

Non-border-Overall and by Grade

Past Year Use

Lifetime Use

Lifetime Use of Selected Substances - Overall							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	6.0%	5.2%	6.7%	0.35%			
ALCOHOL	22.3%	21.0%	23.6%	0.66%			
INHALANTS	9.7%	9.1%	10.3%	0.31%			
MARIJUANA	1.8%	1.5%	2.0%	0.12%			
CIGARETTES	5.1%	4.5%	5.8%	0.33%			
SMOKELESS TOBACCO	1.5%	1.2%	1.7%	0.11%			
BEER	14.2%	13.2%	15.2%	0.51%			
WINE COOLERS	8.1%	7.1%	9.0%	0.48%			
WINE	11.8%	11.1%	12.6%	0.37%			
LIQUOR	5.6%	5.1%	6.1%	0.25%			

Past Year Use of Selected Substances - Overall							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	3.2%	2.8%	3.6%	0.19%			
ALCOHOL	13.7%	12.8%	14.7%	0.48%			
INHALANTS	6.9%	6.4%	7.3%	0.22%			
MARIJUANA	1.2%	1.0%	1.4%	0.09%			
CIGARETTES	2.7%	2.3%	3.0%	0.18%			
SMOKELESS TOBACCO	0.8%	0.7%	0.9%	0.06%			
BEER	7.7%	7.0%	8.3%	0.31%			
WINE COOLERS	4.9%	4.2%	5.5%	0.32%			
WINE	7.1%	6.6%	7.5%	0.22%			
LIQUOR	3.4%	3.1%	3.7%	0.16%			

Lifetime Use of Selected Substances - Grade 4						
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	3.6%	3.1%	4.1%	0.24%		
ALCOHOL	15.8%	14.9%	16.7%	0.46%		
INHALANTS	9.4%	8.7%	10.2%	0.38%		
MARIJUANA	0.7%	0.5%	0.8%	0.08%		
CIGARETTES	2.7%	2.3%	3.2%	0.22%		
SMOKELESS TOBACCO	1.3%	1.1%	1.5%	0.11%		
BEER	10.4%	9.6%	11.1%	0.36%		
WINE COOLERS	3.7%	3.0%	4.3%	0.30%		
WINE	7.4%	6.8%	8.1%	0.32%		
LIQUOR	2.4%	2.1%	2.6%	0.10%		

Past Year Use of Selected Substances - Grade 4						
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	1.8%	1.5%	2.1%	0.15%		
ALCOHOL	9.4%	8.7%	10.1%	0.34%		
INHALANTS	6.4%	5.8%	7.1%	0.32%		
MARIJUANA	0.4%	0.3%	0.5%	0.05%		
CIGARETTES	1.3%	1.1%	1.6%	0.12%		
SMOKELESS TOBACCO	0.6%	0.5%	0.8%	0.07%		
BEER	5.5%	5.0%	5.9%	0.24%		
WINE COOLERS	2.1%	1.7%	2.5%	0.20%		
WINE	4.4%	4.0%	4.7%	0.18%		
LIQUOR	1.4%	1.2%	1.5%	0.09%		

Lifetime Use of Selected Substances - Grade 5						
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	4.9%	4.3%	5.5%	0.30%		
ALCOHOL	20.6%	19.2%	22.0%	0.70%		
INHALANTS	8.4%	7.7%	9.1%	0.37%		
MARIJUANA	1.2%	0.9%	1.4%	0.13%		
CIGARETTES	4.2%	3.7%	4.7%	0.27%		
SMOKELESS TOBACCO	1.2%	0.9%	1.4%	0.12%		
BEER	12.7%	11.6%	13.8%	0.55%		
WINE COOLERS	6.8%	5.8%	7.7%	0.48%		
WINE	10.8%	10.0%	11.5%	0.37%		

Past Year Use of Selected Substances - Grade 5						
CATEGORY	Prevalence	Boun	daries	Standard		
		Lower	Upper	Error		
TOBACCO	2.7%	2.3%	3.0%	0.19%		
ALCOHOL	12.6%	11.6%	13.5%	0.48%		
INHALANTS	6.2%	5.6%	6.7%	0.28%		
MARIJUANA	0.8%	0.6%	0.9%	0.09%		
CIGARETTES	2.1%	1.8%	2.5%	0.19%		
SMOKELESS TOBACCO	0.7%	0.5%	0.8%	0.07%		
BEER	6.8%	6.1%	7.4%	0.33%		
WINE COOLERS	4.2%	3.6%	4.8%	0.31%		
WINE	6.5%	6.0%	7.0%	0.25%		

LIQUOR	4.4%	4.0%	4.9%	0.24%	LIQUOR	2.5%	2.2%	2.8%	0.16%
LIQUUIK	7.770	7.070	7.570	0.27/0	LIQUUN	2.570	2.2/0	2.070	0.10/0

Lifetime Use of Selected Substances - Grade 6							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	9.4%	8.2%	10.6%	0.61%			
ALCOHOL	30.6%	28.3%	32.9%	1.17%			
INHALANTS	11.3%	10.3%	12.3%	0.48%			
MARIJUANA	3.5%	3.0%	4.1%	0.29%			
CIGARETTES	8.5%	7.3%	9.7%	0.59%			
SMOKELESS TOBACCO	1.9%	1.6%	2.3%	0.17%			
BEER	19.7%	17.9%	21.5%	0.90%			
WINE COOLERS	13.9%	12.1%	15.6%	0.87%			
WINE	17.4%	16.0%	18.7%	0.66%			
LIQUOR	10.2%	9.1%	11.4%	0.58%			

Past Year Use of Selected Substances - Grade 6							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	5.1%	4.3%	5.8%	0.36%			
ALCOHOL	19.3%	17.4%	21.1%	0.93%			
INHALANTS	8.1%	7.5%	8.7%	0.31%			
MARIJUANA	2.5%	2.0%	2.9%	0.21%			
CIGARETTES	4.5%	3.8%	5.2%	0.37%			
SMOKELESS TOBACCO	1.1%	0.8%	1.3%	0.11%			
BEER	10.8%	9.7%	12.0%	0.57%			
WINE COOLERS	8.4%	7.1%	9.6%	0.63%			
WINE	10.3%	9.5%	11.1%	0.41%			
LIQUOR	6.4%	5.6%	7.1%	0.39%			

Lifetime Use of Selected Substances - Overall Male						
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	7.3%	6.4%	8.2%	0.44%		
ALCOHOL	25.0%	23.8%	26.3%	0.63%		
INHALANTS	11.0%	10.3%	11.8%	0.36%		
MARIJUANA	2.4%	2.0%	2.7%	0.18%		
CIGARETTES	6.1%	5.3%	6.9%	0.41%		
SMOKELESS TOBACCO	2.2%	1.9%	2.5%	0.16%		
BEER	17.3%	16.2%	18.5%	0.56%		
WINE COOLERS	7.8%	7.0%	8.7%	0.44%		
WINE	13.1%	12.2%	13.9%	0.43%		
LIQUOR	6.5%	5.9%	7.1%	0.31%		

Lifetime Use of Selected Substances - Grade 4 Male							
CATEGORY	Prevalence	Boun	daries	Standard			
		Lower	Upper	Error			
TOBACCO	4.8%	4.2%	5.4%	0.31%			
ALCOHOL	18.9%	17.8%	20.0%	0.55%			
INHALANTS	11.0%	10.0%	12.1%	0.52%			
MARIJUANA	1.0%	0.7%	1.2%	0.13%			
CIGARETTES	3.5%	2.9%	4.2%	0.32%			
SMOKELESS TOBACCO	1.9%	1.6%	2.1%	0.13%			
BEER	13.5%	12.5%	14.6%	0.52%			
WINE COOLERS	3.9%	3.4%	4.4%	0.27%			
WINE	8.7%	7.7%	9.6%	0.49%			
LIQUOR	3.1%	2.7%	3.6%	0.22%			

Lifetime Use of Selected Substances - Grade 5 Male						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	6.1%	5.2%	6.9%	0.44%		
ALCOHOL	23.2%	21.7%	24.7%	0.75%		
INHALANTS	9.9%	8.9%	10.9%	0.48%		
MARIJUANA	1.6%	1.2%	1.9%	0.17%		
CIGARETTES	5.0%	4.3%	5.7%	0.37%		
SMOKELESS TOBACCO	1.8%	1.3%	2.2%	0.22%		
BEER	15.5%	14.3%	16.8%	0.64%		

Past Year Use of Selected Substances - Overall Male				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	3.9%	3.4%	4.4%	0.25%
ALCOHOL	15.2%	14.3%	16.1%	0.47%
INHALANTS	7.8%	7.3%	8.3%	0.26%
MARIJUANA	1.6%	1.3%	1.8%	0.13%
CIGARETTES	3.1%	2.6%	3.6%	0.23%
SMOKELESS TOBACCO	1.2%	1.0%	1.4%	0.09%
BEER	9.4%	8.7%	10.1%	0.35%
WINE COOLERS	4.6%	4.0%	5.2%	0.30%
WINE	7.8%	7.3%	8.3%	0.25%
LIQUOR	3.8%	3.4%	4.2%	0.20%

Past Year Use of Selected Substances - Grade 4 Male				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	2.5%	2.1%	2.9%	0.19%
ALCOHOL	11.1%	10.3%	11.9%	0.42%
INHALANTS	7.7%	7.0%	8.4%	0.37%
MARIJUANA	0.6%	0.4%	0.8%	0.09%
CIGARETTES	1.8%	1.5%	2.1%	0.17%
SMOKELESS TOBACCO	0.9%	0.7%	1.1%	0.11%
BEER	7.1%	6.4%	7.8%	0.35%
WINE COOLERS	2.1%	1.7%	2.4%	0.17%
WINE	5.0%	4.5%	5.6%	0.29%
LIQUOR	1.9%	1.6%	2.1%	0.14%

Past Year Use of Selected Substances - Grade 5 Male					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.4%	2.9%	3.9%	0.25%	
ALCOHOL	14.3%	13.2%	15.4%	0.55%	
INHALANTS	7.2%	6.5%	7.9%	0.36%	
MARIJUANA	1.0%	0.8%	1.2%	0.10%	
CIGARETTES	2.6%	2.1%	3.0%	0.23%	
SMOKELESS TOBACCO	1.1%	0.8%	1.3%	0.12%	
BEER	8.4%	7.5%	9.3%	0.43%	

WINE COOLERS	6.9%	5.9%	7.8%	0.48%
WINE	11.9%	11.0%	12.8%	0.45%
LIQUOR	5.2%	4.6%	5.8%	0.29%

WINE COOLERS	4.3%	3.7%	5.0%	0.32%
WINE	7.4%	6.7%	8.0%	0.32%
LIQUOR	2.9%	2.5%	3.3%	0.21%

Lifetime Use of Selected Substances - Grade 6 Male				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	11.2%	9.7%	12.6%	0.71%
ALCOHOL	33.2%	31.0%	35.4%	1.12%
INHALANTS	12.3%	11.1%	13.4%	0.57%
MARIJUANA	4.6%	3.8%	5.4%	0.39%
CIGARETTES	9.8%	8.5%	11.2%	0.67%
SMOKELESS TOBACCO	2.9%	2.4%	3.5%	0.26%
BEER	23.1%	21.1%	25.1%	0.99%
WINE COOLERS	12.9%	11.1%	14.6%	0.86%
WINE	18.8%	17.5%	20.2%	0.68%
LIQUOR	11.3%	9.9%	12.7%	0.71%

Past Year Use of Selected Substances - Grade 6 Male					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	5.9%	5.0%	6.8%	0.45%	
ALCOHOL	20.3%	18.4%	22.2%	0.95%	
INHALANTS	8.5%	7.7%	9.2%	0.37%	
MARIJUANA	3.2%	2.6%	3.8%	0.30%	
CIGARETTES	5.0%	4.1%	5.9%	0.46%	
SMOKELESS TOBACCO	1.6%	1.3%	1.9%	0.16%	
BEER	12.8%	11.6%	14.0%	0.61%	
WINE COOLERS	7.3%	6.1%	8.6%	0.61%	
WINE	11.0%	10.1%	12.0%	0.48%	
LIQUOR	6.8%	5.8%	7.8%	0.49%	

Lifetime Use of S	Lifetime Use of Selected Substances - Overall Female					
CATEGORY	Prevalence	Bound	aries	Standard		
		Lower	Upper	Error		
TOBACCO	4.7%	4.0%	5.3%	0.31%		
ALCOHOL	19.7%	18.2%	21.2%	0.74%		
INHALANTS	8.4%	7.7%	9.2%	0.37%		
MARIJUANA	1.2%	1.1%	1.4%	0.10%		
CIGARETTES	4.2%	3.6%	4.8%	0.29%		
SMOKELESS TOBACCO	0.8%	0.6%	0.9%	0.07%		
BEER	11.2%	10.2%	12.2%	0.49%		
WINE COOLERS	8.3%	7.2%	9.4%	0.56%		
WINE	10.6%	9.9%	11.4%	0.37%		
LIQUOR	4.8%	4.3%	5.3%	0.25%		

_				
Lifetime Use of S	elected Substance	es - Grade 4	Female	
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	2.5%	2.0%	3.0%	0.23%
ALCOHOL	12.9%	11.8%	13.9%	0.51%
INHALANTS	7.9%	6.9%	8.9%	0.50%
MARIJUANA	0.4%	0.3%	0.5%	0.06%
CIGARETTES	2.0%	1.6%	2.3%	0.20%
SMOKELESS TOBACCO	0.7%	0.5%	1.0%	0.12%
BEER	7.4%	6.6%	8.1%	0.37%
WINE COOLERS	3.4%	2.6%	4.3%	0.42%
WINE	6.2%	5.7%	6.8%	0.27%
LIQUOR	1.6%	1.3%	1.9%	0.16%

Lifetime Use of Selected Substances - Grade 5 Female				
CATEGORY	Prevalence	Bound	Boundaries	
		Lower	Upper	Error
TOBACCO	3.8%	3.2%	4.3%	0.26%
ALCOHOL	18.1%	16.5%	19.7%	0.80%
INHALANTS	7.0%	6.0%	7.9%	0.48%
MARIJUANA	0.8%	0.6%	1.0%	0.11%
CIGARETTES	3.4%	2.9%	3.9%	0.26%
SMOKELESS TOBACCO	0.6%	0.4%	0.8%	0.09%
BEER	9.9%	8.8%	11.0%	0.56%

Past Year Use of Selected Substances - Overall Female					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	2.5%	2.1%	2.8%	0.18%	
ALCOHOL	12.3%	11.2%	13.4%	0.53%	
INHALANTS	6.1%	5.5%	6.6%	0.26%	
MARIJUANA	0.9%	0.7%	1.0%	0.07%	
CIGARETTES	2.2%	1.9%	2.5%	0.16%	
SMOKELESS TOBACCO	0.4%	0.3%	0.5%	0.05%	
BEER	6.0%	5.4%	6.6%	0.29%	
WINE COOLERS	5.2%	4.4%	5.9%	0.37%	
WINE	6.4%	5.9%	6.8%	0.24%	
LIQUOR	3.0%	2.6%	3.3%	0.18%	

Past Year Use of Selected Substances - Grade 4 Female				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	1.2%	0.9%	1.5%	0.16%
ALCOHOL	7.8%	7.0%	8.6%	0.39%
INHALANTS	5.2%	4.5%	5.9%	0.36%
MARIJUANA	0.2%	0.1%	0.3%	0.04%
CIGARETTES	0.9%	0.7%	1.2%	0.12%
SMOKELESS TOBACCO	0.3%	0.2%	0.5%	0.07%
BEER	3.9%	3.4%	4.5%	0.28%
WINE COOLERS	2.1%	1.6%	2.7%	0.28%
WINE	3.8%	3.4%	4.2%	0.20%
LIQUOR	0.9%	0.7%	1.1%	0.11%

Past Year Use of Selected Substances - Grade 5 Female					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	1.9%	1.5%	2.4%	0.22%	
ALCOHOL	10.9%	9.8%	11.9%	0.52%	
INHALANTS	5.2%	4.5%	5.9%	0.36%	
MARIJUANA	0.5%	0.3%	0.7%	0.09%	
CIGARETTES	1.7%	1.2%	2.1%	0.23%	
SMOKELESS TOBACCO	0.3%	0.2%	0.4%	0.06%	
BEER	5.1%	4.5%	5.7%	0.31%	

WINE COOLERS	6.7%	5.5%	7.9%	0.59%
WINE	9.7%	8.8%	10.5%	0.42%
LIQUOR	3.7%	3.2%	4.2%	0.26%

Lifetime Use of Selected Substances -Grade 6 Female						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	7.8%	6.5%	9.0%	0.60%		
ALCOHOL	28.2%	25.6%	30.7%	1.29%		
INHALANTS	10.4%	9.3%	11.5%	0.56%		
MARIJUANA	2.5%	2.0%	3.0%	0.25%		
CIGARETTES	7.3%	6.1%	8.5%	0.60%		
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.12%		
BEER	16.5%	14.7%	18.3%	0.90%		
WINE COOLERS	14.9%	12.9%	16.8%	0.97%		
WINE	16.0%	14.5%	17.4%	0.72%		
LIQUOR	9.2%	8.0%	10.3%	0.57%		

WINE COOLERS	4.0%	3.3%	4.8%	0.37%
WINE	5.7%	5.0%	6.3%	0.32%
LIQUOR	2.1%	1.8%	2.4%	0.16%

Past Year Use of Selected Substances - Grade 6 Female						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	4.3%	3.6%	5.0%	0.35%		
ALCOHOL	18.3%	16.3%	20.3%	0.99%		
INHALANTS	7.8%	7.0%	8.6%	0.42%		
MARIJUANA	1.8%	1.5%	2.2%	0.18%		
CIGARETTES	4.0%	3.3%	4.7%	0.35%		
SMOKELESS TOBACCO	0.6%	0.4%	0.8%	0.10%		
BEER	9.0%	7.8%	10.1%	0.58%		
WINE COOLERS	9.3%	7.9%	10.7%	0.69%		
WINE	9.6%	8.6%	10.6%	0.48%		
LIQUOR	6.0%	5.1%	6.8%	0.42%		

Lifetime Use of Selected Substances - Overall White						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	5.0%	4.2%	5.9%	0.43%		
ALCOHOL	19.5%	17.8%	21.1%	0.84%		
INHALANTS	8.2%	7.4%	8.9%	0.38%		
MARIJUANA	1.0%	0.8%	1.3%	0.13%		
CIGARETTES	3.9%	3.1%	4.7%	0.38%		
SMOKELESS TOBACCO	1.9%	1.5%	2.2%	0.17%		
BEER	12.1%	10.9%	13.4%	0.63%		
WINE COOLERS	5.5%	4.6%	6.4%	0.46%		
WINE	12.0%	10.9%	13.0%	0.52%		
LIQUOR	5.1%	4.5%	5.7%	0.30%		

	Selected Substance		Ct	
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.6%	2.2%	3.0%	0.21%
ALCOHOL	12.2%	11.1%	13.2%	0.51%
INHALANTS	5.7%	5.1%	6.4%	0.32%
MARIJUANA	0.7%	0.5%	0.8%	0.08%
CIGARETTES	1.9%	1.6%	2.3%	0.18%
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.10%
BEER	6.4%	5.7%	7.0%	0.34%
WINE COOLERS	3.2%	2.7%	3.8%	0.28%
WINE	7.2%	6.6%	7.8%	0.30%
LIQUOR	3.2%	2.9%	3.6%	0.17%

Lifetime Use of Selected Substances - Grade 4 White					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.4%	2.6%	4.3%	0.42%	
ALCOHOL	14.7%	13.2%	16.2%	0.76%	
INHALANTS	9.3%	8.0%	10.6%	0.66%	
MARIJUANA	0.4%	0.3%	0.6%	0.09%	
CIGARETTES	2.2%	1.4%	3.0%	0.39%	
SMOKELESS TOBACCO	1.6%	1.2%	2.0%	0.21%	
BEER	9.5%	8.2%	10.8%	0.65%	
WINE COOLERS	2.3%	1.7%	2.9%	0.30%	
WINE	8.3%	7.3%	9.3%	0.50%	
LIQUOR	2.2%	1.6%	2.7%	0.25%	

Past Year Use of Selected Substances - Grade 4 White						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	1.5%	1.1%	2.0%	0.23%		
ALCOHOL	8.7%	7.7%	9.7%	0.50%		
INHALANTS	6.4%	5.4%	7.4%	0.49%		
MARIJUANA	0.3%	0.1%	0.4%	0.08%		
CIGARETTES	0.9%	0.5%	1.3%	0.20%		
SMOKELESS TOBACCO	0.7%	0.4%	1.0%	0.13%		
BEER	4.7%	4.0%	5.4%	0.35%		
WINE COOLERS	1.1%	0.7%	1.6%	0.23%		
WINE	5.0%	4.2%	5.7%	0.37%		
LIQUOR	1.3%	1.0%	1.6%	0.16%		

Lifetime Use of Selected Substances - Grade 5 White						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	4.1%	3.1%	5.1%	0.50%		
ALCOHOL	17.9%	16.3%	19.5%	0.81%		
INHALANTS	6.5%	5.7%	7.3%	0.41%		
MARIJUANA	0.6%	0.4%	0.8%	0.09%		
CIGARETTES	3.2%	2.4%	4.0%	0.41%		
SMOKELESS TOBACCO	1.4%	0.9%	1.9%	0.24%		
BEER	10.9%	9.5%	12.2%	0.66%		

Past Year Use of Selected Substances - Grade 5 White					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	2.3%	1.7%	2.8%	0.27%	
ALCOHOL	10.9%	10.0%	11.9%	0.48%	
INHALANTS	4.8%	4.1%	5.5%	0.35%	
MARIJUANA	0.4%	0.3%	0.5%	0.06%	
CIGARETTES	1.6%	1.1%	2.0%	0.22%	
SMOKELESS TOBACCO	0.9%	0.6%	1.1%	0.14%	
BEER	5.4%	4.6%	6.3%	0.43%	

WINE COOLERS	4.5%	3.4%	5.5%	0.53%
WINE	10.8%	9.8%	11.8%	0.52%
LIQUOR	3.8%	3.3%	4.3%	0.26%

WINE COOLERS	2.8%	2.1%	3.6%	0.37%
WINE	6.6%	5.9%	7.1%	0.30%
LIQUOR	2.4%	1.9%	2.8%	0.22%

Lifetime Use of Selected Substances - Grade 6 White					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	7.5%	6.3%	8.7%	0.60%	
ALCOHOL	25.5%	22.6%	28.3%	1.44%	
INHALANTS	8.7%	7.5%	9.8%	0.56%	
MARIJUANA	2.0%	1.4%	2.5%	0.28%	
CIGARETTES	6.3%	5.2%	7.4%	0.55%	
SMOKELESS TOBACCO	2.5%	2.0%	3.0%	0.24%	
BEER	15.9%	13.9%	17.9%	0.98%	
WINE COOLERS	9.5%	7.9%	11.0%	0.79%	
WINE	16.5%	14.7%	18.3%	0.90%	
LIQUOR	9.1%	7.8%	10.4%	0.64%	

Past Year Use of Selected Substances - Grade 6 White					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	4.0%	3.5%	4.5%	0.27%	
ALCOHOL	16.6%	14.8%	18.5%	0.94%	
INHALANTS	6.1%	5.2%	7.0%	0.45%	
MARIJUANA	1.3%	0.9%	1.7%	0.19%	
CIGARETTES	3.2%	2.7%	3.8%	0.27%	
SMOKELESS TOBACCO	1.3%	1.0%	1.6%	0.17%	
BEER	8.8%	7.6%	9.9%	0.57%	
WINE COOLERS	5.6%	4.7%	6.5%	0.46%	
WINE	10.0%	8.9%	11.0%	0.54%	
LIQUOR	5.9%	5.1%	6.6%	0.37%	

Lifetime Use of Selected Substances - Overall African/American						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	7.0%	5.3%	8.7%	0.85%		
ALCOHOL	26.7%	24.5%	28.9%	1.08%		
INHALANTS	9.0%	7.7%	10.4%	0.66%		
MARIJUANA	2.1%	1.6%	2.5%	0.22%		
CIGARETTES	6.1%	4.6%	7.6%	0.73%		
SMOKELESS TOBACCO	1.6%	1.1%	2.1%	0.26%		
BEER	15.5%	13.6%	17.4%	0.97%		
WINE COOLERS	14.8%	12.5%	17.1%	1.15%		
WINE	11.6%	10.6%	12.5%	0.47%		
LIQUOR	6.0%	4.7%	7.3%	0.64%		

Standard	CATEGORY	Prevalence	Bound	daries	Standard	
Error			Lower	Upper	Error	
0.85%	TOBACCO	3.4%	2.4%	4.4%	0.49%	
1.08%	ALCOHOL	15.2%	13.3%	17.1%	0.94%	
0.66%	INHALANTS	6.1%	5.1%	7.1%	0.49%	
0.22%	MARIJUANA	1.2%	0.9%	1.6%	0.17%	
0.73%	CIGARETTES	2.9%	2.1%	3.8%	0.42%	
0.26%	SMOKELESS TOBACCO	0.7%	0.4%	1.0%	0.16%	
0.97%	BEER	7.4%	6.3%	8.6%	0.59%	
1.15%	WINE COOLERS	8.2%	6.7%	9.8%	0.77%	
0.47%	WINE	6.3%	5.8%	6.9%	0.28%	
0.64%	LIQUOR	3.3%	2.5%	4.0%	0.37%	

Lifetime Use of Selected Substances - Grade 4 African/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.0%	3.6%	6.3%	0.66%	
ALCOHOL	19.4%	17.5%	21.3%	0.96%	
INHALANTS	8.4%	7.1%	9.6%	0.62%	
MARIJUANA	0.8%	0.4%	1.1%	0.19%	
CIGARETTES	4.1%	2.9%	5.2%	0.59%	
SMOKELESS TOBACCO	1.3%	0.9%	1.8%	0.23%	
BEER	12.1%	10.5%	13.7%	0.81%	
WINE COOLERS	8.6%	6.6%	10.6%	1.02%	
WINE	7.2%	6.2%	8.2%	0.49%	
LIQUOR	2.8%	2.1%	3.5%	0.37%	

Past Year Use of Selected Substances - Grade 4 African/American						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	2.7%	1.9%	3.5%	0.41%		
ALCOHOL	11.5%	9.7%	13.3%	0.90%		
INHALANTS	5.5%	4.7%	6.3%	0.40%		
MARIJUANA	0.3%	0.0%	0.6%	0.16%		
CIGARETTES	2.1%	1.4%	2.9%	0.38%		
SMOKELESS TOBACCO	0.6%	0.3%	1.0%	0.19%		
BEER	6.4%	5.1%	7.7%	0.65%		
WINE COOLERS	5.0%	3.7%	6.4%	0.69%		
WINE	4.2%	3.7%	4.8%	0.28%		
LIQUOR	1.5%	0.9%	2.0%	0.28%		

Lifetime Use of Selected Substances - Grade 5 African/American				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	5.8%	5.0%	6.6%	0.38%
ALCOHOL	24.6%	22.6%	26.6%	0.99%
INHALANTS	7.7%	6.2%	9.3%	0.77%
MARIJUANA	1.5%	1.1%	1.9%	0.20%
CIGARETTES	4.7%	3.9%	5.4%	0.38%
SMOKELESS TOBACCO	1.6%	0.9%	2.2%	0.32%
BEER	14.1%	12.9%	15.3%	0.60%

Past Year Use of Selected Substances - Grade 5 African/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.1%	2.4%	3.7%	0.32%	
ALCOHOL	14.3%	13.1%	15.5%	0.60%	
INHALANTS	5.6%	4.2%	7.0%	0.71%	
MARIJUANA	1.1%	0.7%	1.4%	0.16%	
CIGARETTES	2.5%	1.9%	3.0%	0.27%	
SMOKELESS TOBACCO	0.9%	0.5%	1.2%	0.19%	
BEER	6.8%	6.0%	7.5%	0.38%	

WINE COOLERS	13.0%	11.1%	14.9%	0.93%
WINE	9.9%	8.6%	11.2%	0.63%
LIQUOR	4.3%	3.2%	5.4%	0.54%

Lifetime Use of Selected Substances - Grade 6 African/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	10.6%	7.0%	14.2%	1.80%	
ALCOHOL	36.9%	32.0%	41.7%	2.42%	
INHALANTS	11.2%	8.4%	13.9%	1.37%	
MARIJUANA	4.1%	2.8%	5.3%	0.62%	
CIGARETTES	9.9%	6.6%	13.2%	1.65%	
SMOKELESS TOBACCO	2.0%	1.2%	2.9%	0.43%	
BEER	20.7%	16.0%	25.4%	2.35%	
WINE COOLERS	23.3%	18.9%	27.7%	2.19%	
WINE	18.1%	15.6%	20.6%	1.26%	
LIQUOR	11.2%	8.5%	14.0%	1.37%	

WINE COOLERS	7.8%	6.9%	8.7%	0.45%
WINE	5.8%	4.9%	6.6%	0.43%
LIQUOR	2.0%	1.2%	2.8%	0.39%

Past Year Use of Select	ed Substances - Gra	ade 6 Afric	an/America	an	
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	4.6%	2.5%	6.7%	1.05%	
ALCOHOL	20.3%	15.1%	25.4%	2.56%	
INHALANTS	7.3%	5.4%	9.1%	0.94%	
MARIJUANA	2.5%	1.6%	3.5%	0.47%	
CIGARETTES	4.4%	2.4%	6.3%	0.99%	
SMOKELESS TOBACCO	0.6%	0.2%	1.1%	0.22%	
BEER	9.2%	6.6%	11.9%	1.33%	
WINE COOLERS	12.1%	8.3%	15.9%	1.91%	
WINE	9.2%	7.5%	10.9%	0.85%	
LIQUOR	6.6%	4.9%	8.3%	0.86%	

Lifetime Use of Selected Substances - Overall Mexican/American				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	6.6%	5.9%	7.2%	0.32%
ALCOHOL	24.3%	23.1%	25.5%	0.59%
INHALANTS	11.1%	10.5%	11.7%	0.32%
MARIJUANA	2.5%	2.1%	2.9%	0.19%
CIGARETTES	6.0%	5.4%	6.6%	0.31%
SMOKELESS TOBACCO	1.1%	1.0%	1.2%	0.07%
BEER	16.3%	15.4%	17.3%	0.47%
WINE COOLERS	8.8%	8.0%	9.6%	0.38%
WINE	12.2%	11.2%	13.1%	0.45%
LIQUOR	6.3%	5.6%	6.9%	0.31%

Lifetime Use of Selecte	d Substances - Gra	de 4 Mexi	can/Ameri	can
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	3.4%	2.9%	3.9%	0.25%
ALCOHOL	16.4%	15.1%	17.6%	0.62%
INHALANTS	9.8%	8.9%	10.7%	0.46%
MARIJUANA	0.9%	0.7%	1.1%	0.11%
CIGARETTES	2.8%	2.3%	3.2%	0.24%
SMOKELESS TOBACCO	0.9%	0.7%	1.1%	0.12%
BEER	11.2%	10.0%	12.3%	0.58%
WINE COOLERS	3.5%	3.0%	4.0%	0.27%
WINE	7.0%	6.0%	8.0%	0.49%
LIQUOR	2.5%	2.1%	2.9%	0.19%

Lifetime Use of Selected Substances - Grade 5 Mexican/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.4%	4.7%	6.0%	0.31%	
ALCOHOL	22.3%	20.5%	24.0%	0.88%	
INHALANTS	9.9%	8.9%	10.8%	0.47%	
MARIJUANA	1.6%	1.2%	2.0%	0.22%	
CIGARETTES	4.9%	4.3%	5.5%	0.30%	
SMOKELESS TOBACCO	0.8%	0.6%	1.1%	0.11%	
BEER	14.4%	13.1%	15.8%	0.67%	

Past Year Use of Selected Substances - Overall Mexican/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.8%	3.4%	4.2%	0.20%	
ALCOHOL	15.4%	14.4%	16.5%	0.52%	
INHALANTS	8.1%	7.6%	8.7%	0.26%	
MARIJUANA	1.8%	1.5%	2.1%	0.16%	
CIGARETTES	3.4%	3.0%	3.8%	0.20%	
SMOKELESS TOBACCO	0.7%	0.5%	0.7%	0.05%	
BEER	9.5%	8.8%	10.1%	0.34%	
WINE COOLERS	5.6%	4.9%	6.3%	0.34%	
WINE	7.4%	6.7%	8.1%	0.33%	
LIQUOR	3.8%	3.4%	4.2%	0.22%	

Past Year Use of Selected Substances - Grade 4 Mexican/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	1.8%	1.4%	2.2%	0.18%	
ALCOHOL	9.9%	9.2%	10.7%	0.38%	
INHALANTS	6.9%	6.0%	7.7%	0.42%	
MARIJUANA	0.6%	0.5%	0.8%	0.09%	
CIGARETTES	1.4%	1.1%	1.8%	0.16%	
SMOKELESS TOBACCO	0.5%	0.3%	0.7%	0.10%	
BEER	6.2%	5.4%	7.0%	0.38%	
WINE COOLERS	2.2%	1.8%	2.6%	0.21%	
WINE	4.2%	3.6%	4.7%	0.29%	
LIQUOR	1.5%	1.2%	1.8%	0.17%	

Past Year Use of Selected Substances - Grade 5 Mexican/American					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	2.9%	2.4%	3.5%	0.27%	
ALCOHOL	14.0%	12.7%	15.3%	0.64%	
INHALANTS	7.3%	6.5%	8.0%	0.38%	
MARIJUANA	1.0%	0.7%	1.3%	0.15%	
CIGARETTES	2.6%	2.1%	3.1%	0.25%	
SMOKELESS TOBACCO	0.5%	0.3%	0.6%	0.07%	
BEER	8.4%	7.5%	9.3%	0.46%	

WINE COOLERS	7.1%	6.3%	8.0%	0.42%
WINE	11.1%	9.7%	12.4%	0.66%
LIQUOR	5.0%	4.2%	5.8%	0.42%

Lifetime Use of Selected Substances - Grade 6 Mexican/American					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	11.0%	9.8%	12.2%	0.60%	
ALCOHOL	34.2%	32.5%	35.9%	0.83%	
INHALANTS	13.7%	12.6%	14.8%	0.56%	
MARIJUANA	5.1%	4.3%	5.9%	0.41%	
CIGARETTES	10.4%	9.1%	11.7%	0.63%	
SMOKELESS TOBACCO	1.6%	1.3%	1.8%	0.13%	
BEER	23.3%	21.8%	24.8%	0.74%	
WINE COOLERS	15.8%	14.4%	17.1%	0.69%	
WINE	18.3%	17.0%	19.6%	0.67%	
LIQUOR	11.3%	9.9%	12.5%	0.65%	

WINE COOLERS	4.4%	3.8%	5.0%	0.30%
WINE	6.8%	5.7%	7.8%	0.51%
LIQUOR	2.8%	2.3%	3.4%	0.29%

Past Year Use of Selected Substances - Grade 6 Mexican/American					
Past Year Use of Select	eu Substances - Gra	ide o Mexi	canyameri	LdII	
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	6.6%	5.7%	7.5%	0.45%	
ALCOHOL	22.4%	20.6%	24.2%	0.89%	
INHALANTS	10.3%	9.5%	11.1%	0.42%	
MARIJUANA	3.8%	3.0%	4.5%	0.36%	
CIGARETTES	6.1%	5.2%	7.0%	0.45%	
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.10%	
BEER	13.8%	12.6%	15.0%	0.60%	
WINE COOLERS	10.2%	8.9%	11.6%	0.68%	
WINE	11.2%	10.2%	12.3%	0.51%	
LIQUOR	7.1%	6.1%	8.1%	0.51%	

Lifetime Use of S	Lifetime Use of Selected Substances - Overall Border				
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.3%	4.6%	5.9%	0.33%	
ALCOHOL	22.2%	20.9%	23.6%	0.67%	
INHALANTS	10.7%	9.6%	11.8%	0.55%	
MARIJUANA	1.9%	1.5%	2.3%	0.22%	
CIGARETTES	4.8%	4.1%	5.5%	0.34%	
SMOKELESS TOBACCO	0.9%	0.7%	1.1%	0.08%	
BEER	14.7%	13.8%	15.7%	0.48%	
WINE COOLERS	8.3%	7.0%	9.6%	0.66%	
WINE	11.2%	10.2%	12.2%	0.51%	
LIQUOR	5.0%	4.0%	6.0%	0.51%	

Lifetime Use of Selected Substances - Grade 4 Border						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	3.0%	2.7%	3.4%	0.17%		
ALCOHOL	16.3%	14.7%	17.8%	0.79%		
INHALANTS	10.2%	9.0%	11.5%	0.64%		
MARIJUANA	0.8%	0.6%	1.0%	0.10%		
CIGARETTES	2.6%	2.2%	3.0%	0.20%		
SMOKELESS TOBACCO	0.7%	0.4%	1.1%	0.18%		
BEER	10.6%	9.0%	12.2%	0.78%		
WINE COOLERS	3.9%	3.0%	4.8%	0.47%		
WINE	7.1%	5.8%	8.3%	0.63%		
LIQUOR	2.2%	1.8%	2.6%	0.21%		

Lifetime Use of Selected Substances - Grade 5 Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	4.5%	3.8%	5.2%	0.35%	
ALCOHOL	19.7%	18.4%	21.0%	0.66%	
INHALANTS	9.9%	8.3%	11.4%	0.79%	
MARIJUANA	1.1%	0.6%	1.5%	0.22%	
CIGARETTES	4.2%	3.5%	4.9%	0.36%	
SMOKELESS TOBACCO	0.6%	0.4%	0.7%	0.07%	
BEER	13.2%	12.1%	14.3%	0.54%	

Past Year Use of Selected Substances - Overall Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.0%	2.5%	3.5%	0.24%	
ALCOHOL	14.1%	12.4%	15.8%	0.84%	
INHALANTS	8.0%	7.1%	8.9%	0.45%	
MARIJUANA	1.3%	1.0%	1.7%	0.18%	
CIGARETTES	2.7%	2.2%	3.2%	0.23%	
SMOKELESS TOBACCO	0.5%	0.3%	0.7%	0.10%	
BEER	8.6%	7.5%	9.6%	0.54%	
WINE COOLERS	5.3%	3.9%	6.6%	0.68%	
WINE	6.8%	5.8%	7.8%	0.49%	
LIQUOR	3.2%	2.5%	3.9%	0.36%	

Past Year Use of Selected Substances - Grade 4 Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	1.6%	1.3%	1.9%	0.15%	
ALCOHOL	10.0%	8.9%	11.0%	0.52%	
INHALANTS	7.5%	6.1%	8.9%	0.70%	
MARIJUANA	0.6%	0.4%	0.7%	0.07%	
CIGARETTES	1.3%	1.1%	1.5%	0.10%	
SMOKELESS TOBACCO	0.5%	0.1%	0.8%	0.17%	
BEER	5.9%	4.9%	7.0%	0.53%	
WINE COOLERS	2.3%	1.5%	3.0%	0.37%	
WINE	4.2%	3.6%	4.7%	0.29%	
LIQUOR	1.6%	1.2%	2.0%	0.20%	

Past Year Use of Selected Substances - Grade 5 Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower Upper		Error	
TOBACCO	2.4%	2.1%	2.8%	0.18%	
ALCOHOL	12.4%	11.1%	13.6%	0.64%	
INHALANTS	7.3%	6.1%	8.4%	0.58%	
MARIJUANA	0.7%	0.4%	1.0%	0.16%	
CIGARETTES	2.2%	1.8%	2.6%	0.19%	
SMOKELESS TOBACCO	0.3%	0.2%	0.5%	0.07%	
BEER	7.9%	7.3%	8.5%	0.30%	

WINE COOLERS	6.3%	5.7%	6.9%	0.30%
WINE	9.8%	8.7%	10.9%	0.55%
LIQUOR	4.0%	3.1%	4.9%	0.45%

Lifetime Use of Selected Substances - Grade 6 Border						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	8.4%	7.3%	9.4%	0.54%		
ALCOHOL	30.9%	29.2%	32.6%	0.85%		
INHALANTS	12.0%	10.5%	13.5%	0.74%		
MARIJUANA	3.8%	3.0%	4.5%	0.37%		
CIGARETTES	7.7%	6.5%	9.0%	0.61%		
SMOKELESS TOBACCO	1.4%	1.2%	1.6%	0.09%		
BEER	20.4%	19.0%	21.8%	0.70%		
WINE COOLERS	14.7%	12.2%	17.2%	1.24%		
WINE	16.9%	16.1%	17.8%	0.41%		
LIQUOR	8.8%	6.5%	11.0%	1.12%		

WINE COOLERS	3.7%	2.9%	4.4%	0.38%
WINE	5.9%	5.2%	6.6%	0.35%
LIQUOR	2.5%	1.7%	3.3%	0.39%

Past Year Use of Selected Substances - Grade 6 Border						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	5.0%	4.1%	5.9%	0.45%		
ALCOHOL	20.0%	16.9%	23.2%	1.58%		
INHALANTS	9.2%	7.8%	10.5%	0.66%		
MARIJUANA	2.7%	2.0%	3.5%	0.37%		
CIGARETTES	4.6%	3.7%	5.5%	0.46%		
SMOKELESS TOBACCO	0.8%	0.7%	1.0%	0.09%		
BEER	11.9%	9.7%	14.0%	1.05%		
WINE COOLERS	10.0%	7.3%	12.6%	1.34%		
WINE	10.4%	8.4%	12.5%	1.02%		
LIQUOR	5.6%	3.9%	7.4%	0.88%		

Lifetime Use of Selected Substances - Overall Non-Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	6.1%	5.3%	7.0%	0.42%	
ALCOHOL	22.3%	20.7%	23.9%	0.80%	
INHALANTS	9.5%	8.8%	10.2%	0.36%	
MARIJUANA	1.8%	1.5%	2.1%	0.15%	
CIGARETTES	5.2%	4.4%	6.0%	0.39%	
SMOKELESS TOBACCO	1.6%	1.3%	1.9%	0.14%	
BEER	14.1%	12.9%	15.3%	0.61%	
WINE COOLERS	8.0%	6.9%	9.2%	0.57%	
WINE	12.0%	11.1%	12.8%	0.44%	
LIQUOR	5.8%	5.2%	6.4%	0.28%	

CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.2%	2.7%	3.7%	0.23%
ALCOHOL	13.6%	12.5%	14.8%	0.56%
INHALANTS	6.6%	6.2%	7.1%	0.24%
MARIJUANA	1.2%	1.0%	1.4%	0.10%
CIGARETTES	2.6%	2.2%	3.1%	0.21%
SMOKELESS TOBACCO	0.8%	0.7%	1.0%	0.08%
BEER	7.5%	6.8%	8.1%	0.35%
WINE COOLERS	4.8%	4.1%	5.5%	0.36%
WINE	7.1%	6.6%	7.6%	0.24%
LIQUOR	3.4%	3.1%	3.8%	0.18%

Lifetime Use of Selected Substances - Grade 4 Non-Border					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
ТОВАССО	3.8%	3.2%	4.4%	0.30%	
ALCOHOL	15.7%	14.7%	16.7%	0.52%	
INHALANTS	9.3%	8.4%	10.1%	0.45%	
MARIJUANA	0.7%	0.5%	0.8%	0.09%	
CIGARETTES	2.8%	2.2%	3.3%	0.27%	
SMOKELESS TOBACCO	1.4%	1.2%	1.7%	0.12%	
BEER	10.3%	9.5%	11.1%	0.40%	
WINE COOLERS	3.6%	2.9%	4.3%	0.36%	
WINE	7.5%	6.8%	8.2%	0.37%	
LIQUOR	2.4%	2.1%	2.6%	0.12%	

Lifetime Use of Selected Substances - Grade 4 Non-Border							
CATEGORY	Prevalence	Bound	Standard				
		Lower	Upper	Error			
TOBACCO	1.9%	1.5%	2.2%	0.18%			
ALCOHOL	9.3%	8.5%	10.1%	0.39%			
INHALANTS	6.2%	5.5%	6.8%	0.34%			
MARIJUANA	0.4%	0.3%	0.5%	0.05%			
CIGARETTES	1.3%	1.1%	1.6%	0.15%			
SMOKELESS TOBACCO	0.7%	0.5%	0.8%	0.08%			
BEER	5.3%	4.8%	5.9%	0.27%			
WINE COOLERS	2.1%	1.6%	2.5%	0.24%			
WINE	4.4%	4.0%	4.8%	0.21%			
LIQUOR	1.3%	1.1%	1.5%	0.09%			

Lifetime Use of Selected Substances - Grade 5 Non-Border							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	5.0%	4.3%	5.7%	0.36%			
ALCOHOL	20.8%	19.1%	22.5%	0.83%			
INHALANTS	8.1%	7.3%	8.9%	0.40%			
MARIJUANA	1.2%	0.9%	1.5%	0.14%			
CIGARETTES	4.2%	3.6%	4.8%	0.32%			

Lifetime Use of Selected Substances - Grade 5 Non-Border							
CATEGORY	Prevalence	Boun	daries	Standard			
		Lower	Upper	Error			
TOBACCO	2.7%	2.2%	3.1%	0.23%			
ALCOHOL	12.6%	11.5%	13.8%	0.57%			
INHALANTS	5.9%	5.3%	6.5%	0.31%			
MARIJUANA	0.8%	0.6%	1.0%	0.10%			
CIGARETTES	2.1%	1.7%	2.6%	0.22%			

SMOKELESS TOBACCO	1.3%	1.0%	1.6%	0.15%
BEER	12.6%	11.3%	13.9%	0.66%
WINE COOLERS	6.9%	5.7%	8.1%	0.58%
WINE	11.0%	10.1%	11.8%	0.43%
LIQUOR	4.5%	4.0%	5.1%	0.28%

SMOKELESS TOBACCO	0.8%	0.6%	0.9%	0.09%
BEER	6.5%	5.7%	7.3%	0.38%
WINE COOLERS	4.3%	3.6%	5.0%	0.37%
WINE	6.6%	6.0%	7.2%	0.30%
LIQUOR	2.5%	2.1%	2.8%	0.17%

Lifetime Use of Selected Substances - Grade 6 Non-Border							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	9.7%	8.2%	11.1%	0.74%			
ALCOHOL	30.5%	27.7%	33.4%	1.43%			
INHALANTS	11.2%	10.0%	12.3%	0.58%			
MARIJUANA	3.5%	2.8%	4.2%	0.34%			
CIGARETTES	8.7%	7.3%	10.1%	0.71%			
SMOKELESS TOBACCO	2.1%	1.6%	2.5%	0.21%			
BEER	19.5%	17.3%	21.7%	1.10%			
WINE COOLERS	13.7%	11.6%	15.8%	1.04%			
WINE	17.5%	15.9%	19.1%	0.81%			
LIQUOR	10.5%	9.2%	11.9%	0.66%			

Lifetime Use of Selected Substances - Grade 6 Non-Border							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	5.1%	4.2%	5.9%	0.42%			
ALCOHOL	19.1%	16.9%	21.3%	1.08%			
INHALANTS	7.9%	7.1%	8.6%	0.35%			
MARIJUANA	2.4%	1.9%	2.9%	0.25%			
CIGARETTES	4.5%	3.6%	5.3%	0.43%			
SMOKELESS TOBACCO	1.1%	0.8%	1.4%	0.13%			
BEER	10.6%	9.3%	11.9%	0.66%			
WINE COOLERS	8.0%	6.6%	9.4%	0.70%			
WINE	10.3%	9.3%	11.2%	0.45%			
LIQUOR	6.5%	5.7%	7.4%	0.43%			

Secondary Grades - 2006

Standard Error and Confidence Intervals

All Students-Overall and by Grade

Past 30 Day Use

Lifetime Use

Male-Overall and by Grade

Past 30 Day Use

Lifetime Use

Female-Overall and by Grade

Past 30 Day Use

Lifetime Use

Whites-Overall and by Grade

Past 30 Day Use

Lifetime Use

African American-Overall and by Grade

Past 30 Day Use

Lifetime Use

Mexican American-Overall and by Grade

Past 30 Day Use

Lifetime Use

Border-Overall and by Grade

Past 30 Day Use

Lifetime Use

Non-border-Overall and by Grade

Past 30 Day Use

Lifetime Use

Lifetime Use of Selected Substances - Overall							
CATEGORY	Prevalence	Boun	daries	Standard			
		Lower	Upper	Error			
TOBACCO	35.4%	33.6%	37.1%	0.86%			
CIGARETTES	33.7%	32.1%	35.3%	0.82%			
SMOKELESS TOBACCO	9.9%	9.0%	10.8%	0.46%			
ALCOHOL	65.9%	64.4%	67.4%	0.75%			
INHALANTS	16.6%	16.0%	17.3%	0.34%			
ANY ILLICIT DRUG	28.0%	26.7%	29.3%	0.64%			
MARIJUANA	26.5%	25.2%	27.7%	0.62%			
MARIJUANA ONLY	13.8%	13.0%	14.6%	0.40%			
COCAINE OR CRACK	7.9%	7.1%	8.7%	0.40%			
COCAINE	7.3%	6.5%	8.1%	0.40%			
CRACK	2.4%	2.2%	2.5%	0.09%			
HALLUCINOGENS	4.7%	4.2%	5.2%	0.25%			
UPPERS	5.7%	5.0%	6.4%	0.36%			
DOWNERS	5.9%	5.3%	6.6%	0.34%			
ROHYPNOL	2.6%	2.3%	2.9%	0.15%			
STEROIDS	1.5%	1.4%	1.6%	0.06%			
ECSTASY	5.2%	4.8%	5.7%	0.20%			
HEROIN	1.5%	1.4%	1.6%	0.06%			

Past 30 Day Use of Selected Substances - Overall						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	15.2%	14.2%	16.3%	0.53%		
CIGARETTES	14.0%	13.0%	14.9%	0.48%		
SMOKELESS TOBACCO	3.9%	3.5%	4.3%	0.21%		
ALCOHOL	31.5%	30.2%	32.8%	0.67%		
INHALANTS	5.8%	5.1%	6.4%	0.34%		
ANY ILLICIT DRUG	12.6%	11.9%	13.3%	0.36%		
MARIJUANA	11.0%	10.4%	11.7%	0.33%		
MARIJUANA ONLY	4.3%	3.9%	4.7%	0.19%		
COCAINE OR CRACK	2.9%	2.5%	3.2%	0.17%		
COCAINE	2.6%	2.3%	2.9%	0.16%		
CRACK	0.7%	0.6%	0.8%	0.03%		
HALLUCINOGENS	1.4%	1.3%	1.6%	0.08%		
UPPERS	2.3%	2.0%	2.6%	0.16%		
DOWNERS	2.5%	2.2%	2.8%	0.16%		
ROHYPNOL	1.0%	0.8%	1.1%	0.06%		
STEROIDS	0.6%	0.5%	0.6%	0.03%		
ECSTASY	1.8%	1.6%	2.0%	0.11%		
HEROIN	0.5%	0.5%	0.6%	0.03%		

Lifetime Use of Selected Substances - Grade 7							
CATEGORY	Prevalence	Boundaries		Standard			
		Lower	Upper	Error			
TOBACCO	18.7%	17.1%	20.4%	0.82%			
CIGARETTES	17.7%	16.1%	19.3%	0.80%			
SMOKELESS TOBACCO	4.5%	4.1%	5.0%	0.23%			
ALCOHOL	47.0%	44.8%	49.1%	1.06%			
INHALANTS	18.4%	8.4% 17.2% 19.5%	0.58%				
ANY ILLICIT DRUG	10.9%	9.4%	12.4%	0.76%			
MARIJUANA	9.4%	8.0%	10.8%	0.70%			
MARIJUANA ONLY	5.5%	4.7%	6.4%	0.43%			
COCAINE OR CRACK	3.1%	2.5%	3.6%	0.28%			
COCAINE	2.3%	1.8%	2.8%	0.23%			
CRACK	1.8%	1.5%	2.1%	0.15%			
HALLUCINOGENS	1.2%	0.9%	1.4%	0.13%			
UPPERS	0.9%	0.8%	1.1%	0.09%			
DOWNERS	0.9%	0.8%	1.1%	0.09%			
ROHYPNOL	1.2%	0.9%	1.5%	0.15%			
STEROIDS	1.3%	1.2%	1.5%	0.07%			
ECSTASY	1.5%	1.3%	1.7%	0.11%			

Past 30 Day Use of Selected Substances - Grade 7							
CATEGORY	Prevalence	Boun	daries	Standard			
		Lower	Upper	Error			
TOBACCO	6.1%	5.3%	6.9%	0.40%			
CIGARETTES	5.5%	4.7%	6.2%	0.36%			
SMOKELESS TOBACCO	1.5%	1.2%	1.8%	0.13%			
ALCOHOL	16.2%	14.7%	17.7%	0.75%			
INHALANTS	7.7%	6.8%	8.6%	0.45%			
ANY ILLICIT DRUG	4.8%	3.9%	5.7%	0.45%			
MARIJUANA	3.9%	3.1%	4.7%	0.38%			
MARIJUANA ONLY	1.9%	1.5%	2.3%	0.22%			
COCAINE OR CRACK	1.0%	0.7%	1.2%	0.12%			
COCAINE	0.7%	0.5%	0.9%	0.11%			
CRACK	0.6%	0.4%	0.7%	0.06%			
HALLUCINOGENS	0.5%	0.3%	0.6%	0.07%			
UPPERS	0.5%	0.3%	0.6%	0.06%			
DOWNERS	0.4%	0.3%	0.5%	0.06%			
ROHYPNOL	0.6%	0.4%	0.7%	0.09%			
STEROIDS	0.5%	0.4%	0.6%	0.05%			
ECSTASY	0.5%	0.4%	0.6%	0.05%			

	I	l l		1	1		l		
HEROIN	1 1%	0.9%	1.3%	0.09%	l HEROIN	0.5%	በ 3%	0.6%	0.05%
I LILINOIN	1.1/0	0.570	1.3/0	0.03/0	ITILITOIN	0.5/0	0.5%	0.070	0.03/0

Lifetime Use of Selected Substances - Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	27.4%	25.5%	29.4%	0.97%
CIGARETTES	25.8%	23.9%	27.7%	0.96%
SMOKELESS TOBACCO	7.0%	6.2%	7.8%	0.40%
ALCOHOL	59.7%	57.8%	61.6%	0.97%
INHALANTS	20.1%	19.1%	21.1%	0.50%
ANY ILLICIT DRUG	18.5%	16.8%	20.1%	0.84%
MARIJUANA	17.0%	15.3%	18.6%	0.82%
MARIJUANA ONLY	9.6%	8.6%	10.6%	0.49%
COCAINE OR CRACK	4.9%	4.2%	5.6%	0.34%
COCAINE	4.0%	3.4%	4.6%	0.31%
CRACK	2.4%	2.0%	2.7%	0.16%
HALLUCINOGENS	2.5%	2.1%	2.7%	0.15%
UPPERS	2.5%	2.1%	2.8%	0.17%
DOWNERS	2.3%	2.0%	2.7%	0.19%
ROHYPNOL	2.3%	1.8%	2.7%	0.23%
STEROIDS	1.5%	1.3%	1.7%	0.11%
ECSTASY	3.0%	2.7%	3.3%	0.17%
HEROIN	1.7%	1.5%	1.9%	0.09%

Lifetime U	Jse of Selected Substa	nces - Grad	de 9	
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	36.2%	34.2%	38.1%	0.95%
CIGARETTES	34.3%	32.4%	36.2%	0.96%
SMOKELESS TOBACCO	9.7%	8.7%	10.6%	0.48%
ALCOHOL	67.8%	66.2%	69.4%	0.79%
INHALANTS	18.0%	16.8%	19.2%	0.60%
ANY ILLICIT DRUG	28.1%	26.5%	29.6%	0.77%
MARIJUANA	26.1%	24.6%	27.7%	0.77%
MARIJUANA ONLY	14.0%	12.8%	15.1%	0.58%
COCAINE OR CRACK	7.5%	6.6%	8.3%	0.43%
COCAINE	6.8%	6.0%	7.7%	0.43%
CRACK	2.5%	2.2%	2.8%	0.15%
HALLUCINOGENS	4.3%	3.8%	4.9%	0.27%
UPPERS	5.0%	4.5%	5.6%	0.29%
DOWNERS	5.2%	4.6%	5.8%	0.28%

Past 30 Day Use of Selected Substances - Grade 8					
CATEGORY	Prevalence	Boun	Boundaries		
		Lower	Upper	Error	
TOBACCO	9.1%	8.2%	10.1%	0.47%	
CIGARETTES	8.3%	7.4%	9.1%	0.44%	
SMOKELESS TOBACCO	2.3%	1.9%	2.7%	0.19%	
ALCOHOL	22.9%	21.4%	24.5%	0.77%	
INHALANTS	7.6%	6.6%	8.7%	0.52%	
ANY ILLICIT DRUG	7.8%	6.9%	8.7%	0.45%	
MARIJUANA	6.9%	6.0%	7.7%	0.43%	
MARIJUANA ONLY	3.1%	2.7%	3.6%	0.22%	
COCAINE OR CRACK	1.7%	1.4%	2.0%	0.13%	
COCAINE	1.4%	1.2%	1.7%	0.12%	
CRACK	0.7%	0.6%	0.9%	0.07%	
HALLUCINOGENS	0.7%	0.6%	0.8%	0.05%	
UPPERS	0.9%	0.8%	1.1%	0.09%	
DOWNERS	0.9%	0.7%	1.1%	0.10%	
ROHYPNOL	0.8%	0.7%	1.0%	0.08%	
STEROIDS	0.5%	0.4%	0.6%	0.06%	
ECSTASY	1.0%	0.8%	1.2%	0.10%	
HEROIN	0.5%	0.4%	0.6%	0.05%	

Past 30 Day Use of Selected Substances - Grade 9					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	14.5%	13.5%	15.4%	0.47%	
CIGARETTES	13.0%	12.1%	13.9%	0.44%	
SMOKELESS TOBACCO	3.7%	3.3%	4.2%	0.23%	
ALCOHOL	31.7%	30.3%	33.2%	0.73%	
INHALANTS	6.3%	5.3%	7.3%	0.49%	
ANY ILLICIT DRUG	12.7%	11.8%	13.6%	0.44%	
MARIJUANA	11.0%	10.1%	11.9%	0.45%	
MARIJUANA ONLY	4.6%	4.0%	5.2%	0.32%	
COCAINE OR CRACK	2.9%	2.5%	3.4%	0.22%	
COCAINE	2.6%	2.2%	3.0%	0.20%	
CRACK	0.8%	0.7%	1.0%	0.09%	
HALLUCINOGENS	1.4%	1.2%	1.6%	0.11%	
UPPERS	1.9%	1.6%	2.2%	0.14%	
DOWNERS	2.1%	1.8%	2.4%	0.15%	

ROHYPNOL	3.0%	2.7%	3.4%	0.19%
STEROIDS	1.3%	1.1%	1.5%	0.09%
ECSTASY	5.1%	4.7%	5.5%	0.21%
HEROIN	1.8%	1.5%	2.1%	0.14%

ROHYPNOL	1.2%	1.0%	1.4%	0.11%
STEROIDS	0.5%	0.4%	0.5%	0.05%
ECSTASY	1.9%	1.6%	2.2%	0.13%
HEROIN	0.6%	0.4%	0.7%	0.07%

Lifetime Use of Selected Substances - Grade 10				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	40.4%	38.6%	42.3%	0.94%
CIGARETTES	38.4%	36.6%	40.2%	0.92%
SMOKELESS TOBACCO	11.5%	10.4%	12.6%	0.55%
ALCOHOL	72.7%	71.1%	74.2%	0.77%
INHALANTS	16.1%	15.0%	17.1%	0.52%
ANY ILLICIT DRUG	34.0%	32.4%	35.7%	0.84%
MARIJUANA	32.3%	30.6%	34.0%	0.83%
MARIJUANA ONLY	16.5%	15.3%	17.6%	0.58%
COCAINE OR CRACK	9.4%	8.3%	10.4%	0.52%
COCAINE	8.9%	7.9%	10.0%	0.52%
CRACK	2.5%	2.2%	2.8%	0.15%
HALLUCINOGENS	6.5%	5.8%	7.2%	0.33%
UPPERS	7.6%	6.7%	8.5%	0.46%
DOWNERS	8.0%	7.1%	8.9%	0.44%
ROHYPNOL	3.4%	2.9%	3.9%	0.24%
STEROIDS	1.7%	1.5%	2.0%	0.11%
ECSTASY	6.8%	6.2%	7.4%	0.30%
HEROIN	1.5%	1.2%	1.7%	0.12%

Past 30 Day Use of Selected Substances - Grade 10					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.7%	16.5%	19.0%	0.61%	
CIGARETTES	16.2%	15.0%	17.3%	0.58%	
SMOKELESS TOBACCO	4.4%	4.0%	4.9%	0.22%	
ALCOHOL	36.8%	35.4%	38.3%	0.72%	
INHALANTS	5.0%	4.0%	6.0%	0.49%	
ANY ILLICIT DRUG	15.7%	14.8%	16.6%	0.45%	
MARIJUANA	13.4%	12.5%	14.2%	0.43%	
MARIJUANA ONLY	4.9%	4.4%	5.4%	0.24%	
COCAINE OR CRACK	3.5%	2.9%	4.0%	0.26%	
COCAINE	3.2%	2.7%	3.7%	0.25%	
CRACK	0.7%	0.6%	0.9%	0.06%	
HALLUCINOGENS	2.0%	1.7%	2.4%	0.17%	
UPPERS	3.4%	2.9%	3.9%	0.24%	
DOWNERS	3.8%	3.3%	4.2%	0.24%	
ROHYPNOL	1.2%	0.9%	1.4%	0.10%	
STEROIDS	0.7%	0.5%	0.8%	0.07%	
ECSTASY	2.4%	2.0%	2.7%	0.17%	
HEROIN	0.6%	0.5%	0.7%	0.06%	

Lifetime Use of Selected Substances - Grade 11				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	44.8%	42.8%	46.8%	1.02%
CIGARETTES	43.0%	41.1%	44.9%	0.96%
SMOKELESS TOBACCO	13.5%	12.2%	14.8%	0.65%
ALCOHOL	75.1%	73.6%	76.6%	0.73%
INHALANTS	13.3%	12.2%	14.3%	0.54%
ANY ILLICIT DRUG	39.8%	38.4%	41.1%	0.68%
MARIJUANA	38.2%	36.8%	39.7%	0.70%
MARIJUANA ONLY	19.5%	18.4%	20.6%	0.56%
COCAINE OR CRACK	11.6%	10.3%	12.8%	0.61%
COCAINE	11.3%	10.1%	12.5%	0.61%
CRACK	2.5%	2.2%	2.8%	0.15%

Past 30 Day Use of Selected Substances - Grade 11					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	21.8%	20.3%	23.3%	0.74%	
CIGARETTES	20.1%	18.7%	21.4%	0.68%	
SMOKELESS TOBACCO	5.5%	5.0%	6.1%	0.26%	
ALCOHOL	40.3%	38.7%	41.9%	0.80%	
INHALANTS	3.9%	3.1%	4.6%	0.37%	
ANY ILLICIT DRUG	18.3%	17.1%	19.5%	0.59%	
MARIJUANA	16.2%	15.0%	17.4%	0.58%	
MARIJUANA ONLY	5.9%	5.2%	6.6%	0.35%	
COCAINE OR CRACK	4.3%	3.8%	4.9%	0.28%	
COCAINE	4.2%	3.6%	4.7%	0.28%	
CRACK	0.7%	0.5%	0.8%	0.07%	

HALLUCINOGENS	7.1%	6.0%	8.1%	0.51%
UPPERS	9.8%	8.4%	11.2%	0.69%
DOWNERS	10.4%	9.1%	11.7%	0.65%
ROHYPNOL	3.0%	2.6%	3.5%	0.24%
STEROIDS	1.5%	1.3%	1.7%	0.10%
ECSTASY	7.6%	6.7%	8.6%	0.48%
HEROIN	1.6%	1.4%	1.9%	0.13%

HALLUCINOGENS	2.0%	1.6%	2.5%	0.22%
UPPERS	4.1%	3.4%	4.8%	0.35%
DOWNERS	4.4%	3.7%	5.0%	0.34%
ROHYPNOL	1.1%	0.9%	1.3%	0.11%
STEROIDS	0.7%	0.6%	0.8%	0.06%
ECSTASY	2.5%	1.9%	3.0%	0.25%
HEROIN	0.4%	0.3%	0.5%	0.06%

Lifetime Use of Selected Substances - Grade 12				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	49.9%	47.2%	52.6%	1.34%
CIGARETTES	48.2%	45.5%	50.9%	1.34%
SMOKELESS TOBACCO	15.5%	13.8%	17.2%	0.86%
ALCOHOL	77.9%	76.1%	79.7%	0.90%
INHALANTS	12.1%	11.3%	12.9%	0.41%
ANY ILLICIT DRUG	42.8%	41.0%	44.6%	0.91%
MARIJUANA	41.6%	39.8%	43.5%	0.91%
MARIJUANA ONLY	20.9%	19.8%	21.9%	0.52%
COCAINE OR CRACK	12.9%	11.4%	14.4%	0.75%
COCAINE	12.7%	11.2%	14.2%	0.75%
CRACK	2.5%	2.2%	2.8%	0.17%
HALLUCINOGENS	8.4%	7.5%	9.4%	0.47%
UPPERS	10.2%	8.7%	11.7%	0.75%
DOWNERS	10.9%	9.5%	12.4%	0.72%
ROHYPNOL	2.9%	2.4%	3.3%	0.23%
STEROIDS	1.8%	1.5%	2.1%	0.13%
ECSTASY	8.9%	8.0%	9.8%	0.45%
HEROIN	1.4%	1.2%	1.7%	0.13%

Past 30 Day Use of Selected Substances - Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	26.2%	24.0%	28.3%	1.07%
CIGARETTES	24.7%	22.7%	26.6%	0.98%
SMOKELESS TOBACCO	6.7%	5.6%	7.8%	0.55%
ALCOHOL	46.3%	43.9%	48.7%	1.20%
INHALANTS	3.0%	2.4%	3.5%	0.26%
ANY ILLICIT DRUG	19.2%	18.0%	20.4%	0.60%
MARIJUANA	17.3%	16.2%	18.4%	0.55%
MARIJUANA ONLY	6.2%	5.7%	6.7%	0.26%
COCAINE OR CRACK	4.5%	3.9%	5.0%	0.29%
COCAINE	4.3%	3.8%	4.9%	0.29%
CRACK	0.6%	0.4%	0.7%	0.07%
HALLUCINOGENS	2.4%	2.1%	2.6%	0.14%
UPPERS	3.9%	3.1%	4.6%	0.36%
DOWNERS	4.3%	3.5%	5.0%	0.36%
ROHYPNOL	0.9%	0.7%	1.1%	0.09%
STEROIDS	0.8%	0.6%	1.0%	0.09%
ECSTASY	2.9%	2.4%	3.4%	0.27%
HEROIN	0.5%	0.4%	0.7%	0.08%

Lifetime Use of Selected Substances - Male Overall				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	37.4%	35.4%	39.4%	1.01%
CIGARETTES	34.5%	32.7%	36.3%	0.91%
SMOKELESS TOBACCO	14.9%	13.2%	16.6%	0.85%
ALCOHOL	64.2%	62.6%	65.8%	0.81%
INHALANTS	16.8%	16.1%	17.4%	0.32%
ANY ILLICIT DRUG	30.3%	28.9%	31.7%	0.69%
MARIJUANA	29.1%	27.8%	30.4%	0.67%
MARIJUANA ONLY	15.8%	14.9%	16.7%	0.44%
COCAINE OR CRACK	8.2%	7.3%	9.1%	0.45%
COCAINE	7.7%	6.8%	8.6%	0.45%
CRACK	2.3%	2.1%	2.5%	0.11%
HALLUCINOGENS	5.8%	5.2%	6.5%	0.32%
UPPERS	5.3%	4.5%	6.0%	0.35%
DOWNERS	5.8%	5.1%	6.5%	0.35%
ROHYPNOL	2.5%	2.2%	2.8%	0.15%
STEROIDS	2.2%	2.0%	2.4%	0.11%
ECSTASY	5.2%	4.7%	5.7%	0.25%
HEROIN	1.6%	1.4%	1.7%	0.08%

Lifetime Use of	Selected Substanc	es - Male C	Overall	
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	17.5%	16.1%	18.8%	0.68%
CIGARETTES	15.2%	14.0%	16.3%	0.56%
SMOKELESS TOBACCO	6.3%	5.4%	7.1%	0.43%
ALCOHOL	31.4%	29.9%	33.0%	0.76%
INHALANTS	5.6%	4.8%	6.4%	0.39%
ANY ILLICIT DRUG	14.1%	13.3%	14.8%	0.38%
MARIJUANA	12.5%	11.8%	13.2%	0.35%
MARIJUANA ONLY	5.1%	4.7%	5.6%	0.21%
COCAINE OR CRACK	3.2%	2.7%	3.6%	0.21%
COCAINE	2.9%	2.5%	3.3%	0.20%
CRACK	0.7%	0.6%	0.8%	0.04%
HALLUCINOGENS	1.9%	1.6%	2.1%	0.11%
UPPERS	2.2%	1.8%	2.5%	0.16%
DOWNERS	2.5%	2.2%	2.9%	0.18%
ROHYPNOL	1.0%	0.9%	1.1%	0.07%
STEROIDS	1.0%	0.8%	1.1%	0.06%
ECSTASY	2.0%	1.7%	2.2%	0.12%
HEROIN	0.6%	0.6%	0.7%	0.04%

Lifetime Use of Selected Substances - Male Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	20.7%	18.7%	22.7%	0.99%
CIGARETTES	19.1%	17.2%	20.9%	0.95%
SMOKELESS TOBACCO	6.2%	5.6%	6.9%	0.34%
ALCOHOL	46.8%	44.3%	49.2%	1.23%
INHALANTS	18.3%	17.0%	19.6%	0.64%
ANY ILLICIT DRUG	12.4%	10.5%	14.3%	0.96%
MARIJUANA	11.3%	9.4%	13.1%	0.90%
MARIJUANA ONLY	7.0%	5.8%	8.2%	0.60%
COCAINE OR CRACK	3.0%	2.3%	3.6%	0.31%
COCAINE	2.3%	1.8%	2.8%	0.26%
CRACK	1.7%	1.4%	2.0%	0.16%
HALLUCINOGENS	1.3%	1.0%	1.7%	0.17%
UPPERS	1.0%	0.8%	1.2%	0.11%
DOWNERS	0.9%	0.7%	1.2%	0.12%
ROHYPNOL	1.2%	0.9%	1.5%	0.16%

Lifetime Use of Selected Substances - Male Grade 7				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	6.8%	5.7%	7.8%	0.53%
CIGARETTES	5.8%	4.8%	6.7%	0.46%
SMOKELESS TOBACCO	2.1%	1.7%	2.5%	0.20%
ALCOHOL	16.2%	14.4%	17.9%	0.87%
INHALANTS	7.3%	6.3%	8.4%	0.53%
ANY ILLICIT DRUG	5.4%	4.2%	6.5%	0.56%
MARIJUANA	4.4%	3.5%	5.4%	0.47%
MARIJUANA ONLY	2.4%	1.8%	2.9%	0.27%
COCAINE OR CRACK	0.9%	0.7%	1.2%	0.13%
COCAINE	0.7%	0.5%	1.0%	0.12%
CRACK	0.5%	0.4%	0.7%	0.08%
HALLUCINOGENS	0.5%	0.3%	0.7%	0.10%
UPPERS	0.4%	0.3%	0.6%	0.08%
DOWNERS	0.4%	0.2%	0.5%	0.07%
ROHYPNOL	0.6%	0.4%	0.7%	0.09%

STEROIDS	1.8%	1.6%	2.1%	0.13%
ECSTASY	1.3%	1.1%	1.6%	0.11%
HEROIN	1.2%	0.9%	1.5%	0.14%

STEROIDS	0.7%	0.6%	0.9%	0.07%
ECSTASY	0.6%	0.4%	0.7%	0.08%
HEROIN	0.6%	0.4%	0.7%	0.08%

Lifetime Use of Selected Substances - Male Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	28.3%	26.1%	30.6%	1.12%
CIGARETTES	25.9%	23.8%	28.0%	1.05%
SMOKELESS TOBACCO	9.4%	8.0%	10.7%	0.69%
ALCOHOL	58.0%	55.9%	60.1%	1.04%
INHALANTS	18.9%	18.0%	19.8%	0.47%
ANY ILLICIT DRUG	20.0%	18.0%	21.9%	0.99%
MARIJUANA	18.9%	16.9%	20.9%	1.01%
MARIJUANA ONLY	11.5%	10.1%	12.8%	0.66%
COCAINE OR CRACK	4.5%	3.8%	5.2%	0.34%
COCAINE	3.8%	3.1%	4.5%	0.33%
CRACK	2.0%	1.6%	2.3%	0.17%
HALLUCINOGENS	2.9%	2.5%	3.3%	0.19%
UPPERS	2.0%	1.5%	2.5%	0.23%
DOWNERS	2.0%	1.5%	2.5%	0.25%
ROHYPNOL	2.2%	1.6%	2.7%	0.27%
STEROIDS	1.8%	1.5%	2.2%	0.18%
ECSTASY	2.6%	2.2%	3.0%	0.21%
HEROIN	1.5%	1.3%	1.7%	0.12%

Lifetime Use of Selected Substances - Male Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	9.8%	8.5%	11.1%	0.63%
CIGARETTES	8.4%	7.4%	9.5%	0.52%
SMOKELESS TOBACCO	3.2%	2.4%	4.0%	0.40%
ALCOHOL	22.2%	20.5%	23.8%	0.82%
INHALANTS	7.1%	5.9%	8.3%	0.60%
ANY ILLICIT DRUG	8.2%	7.2%	9.1%	0.48%
MARIJUANA	7.3%	6.4%	8.1%	0.44%
MARIJUANA ONLY	3.6%	3.1%	4.1%	0.24%
COCAINE OR CRACK	1.7%	1.4%	2.1%	0.17%
COCAINE	1.4%	1.1%	1.8%	0.17%
CRACK	0.6%	0.5%	0.8%	0.07%
HALLUCINOGENS	0.9%	0.7%	1.0%	0.07%
UPPERS	0.8%	0.6%	1.0%	0.10%
DOWNERS	0.8%	0.5%	1.0%	0.14%
ROHYPNOL	0.8%	0.6%	1.0%	0.10%
STEROIDS	0.8%	0.6%	1.0%	0.10%
ECSTASY	0.9%	0.7%	1.2%	0.12%
HEROIN	0.6%	0.4%	0.7%	0.06%

Lifetime Use of Selected Substances - Male Grade 9				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	37.7%	35.5%	40.0%	1.11%
CIGARETTES	34.5%	32.3%	36.8%	1.12%
SMOKELESS TOBACCO	13.7%	11.9%	15.5%	0.90%
ALCOHOL	65.6%	64.0%	67.1%	0.77%
INHALANTS	17.0%	15.6%	18.4%	0.72%
ANY ILLICIT DRUG	30.8%	28.9%	32.7%	0.94%
MARIJUANA	29.4%	27.5%	31.2%	0.93%
MARIJUANA ONLY	16.6%	15.3%	18.0%	0.67%
COCAINE OR CRACK	7.5%	6.4%	8.6%	0.55%
COCAINE	6.9%	5.9%	8.0%	0.53%
CRACK	2.5%	2.0%	2.9%	0.23%
HALLUCINOGENS	5.3%	4.6%	6.0%	0.35%

Lifetime Use of Selected Substances - Male Grade 9					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	16.0%	14.7%	17.2%	0.63%	
CIGARETTES	13.4%	12.3%	14.4%	0.52%	
SMOKELESS TOBACCO	5.6%	4.6%	6.6%	0.50%	
ALCOHOL	30.9%	29.2%	32.6%	0.84%	
INHALANTS	5.7%	4.6%	6.7%	0.53%	
ANY ILLICIT DRUG	13.9%	12.7%	15.1%	0.60%	
MARIJUANA	12.2%	11.0%	13.4%	0.61%	
MARIJUANA ONLY	5.3%	4.5%	6.2%	0.42%	
COCAINE OR CRACK	3.1%	2.5%	3.8%	0.31%	
COCAINE	2.8%	2.3%	3.4%	0.28%	
CRACK	0.9%	0.7%	1.1%	0.11%	
HALLUCINOGENS	1.7%	1.3%	2.1%	0.20%	

_				
UPPERS	4.4%	3.7%	5.0%	0.32%
DOWNERS	4.9%	4.2%	5.6%	0.36%
ROHYPNOL	3.0%	2.5%	3.5%	0.25%
STEROIDS	1.7%	1.4%	2.1%	0.16%
ECSTASY	4.9%	4.3%	5.4%	0.27%
HEROIN	1.8%	1.4%	2.1%	0.16%

UPPERS	1.7%	1.4%	2.1%	0.18%
DOWNERS	2.1%	1.6%	2.5%	0.21%
ROHYPNOL	1.3%	1.0%	1.6%	0.16%
STEROIDS	0.7%	0.5%	0.8%	0.07%
ECSTASY	2.0%	1.6%	2.3%	0.18%
HEROIN	0.6%	0.5%	0.8%	0.08%

Lifetime Use of Selected Substances - Male Grade 10					
CATEGORY	Prevalence		daries	Standard	
		Lower	Upper	Error	
TOBACCO	42.3%	40.1%	44.4%	1.08%	
CIGARETTES	38.4%	36.4%	40.5%	1.01%	
SMOKELESS TOBACCO	17.5%	15.6%	19.4%	0.95%	
ALCOHOL	70.3%	68.7%	71.9%	0.81%	
INHALANTS	15.8%	14.8%	16.8%	0.48%	
ANY ILLICIT DRUG	36.3%	34.4%	38.1%	0.92%	
MARIJUANA	34.8%	33.0%	36.5%	0.87%	
MARIJUANA ONLY	18.8%	17.5%	20.1%	0.66%	
COCAINE OR CRACK	9.5%	8.5%	10.6%	0.53%	
COCAINE	9.1%	8.0%	10.1%	0.53%	
CRACK	2.4%	2.0%	2.8%	0.18%	
HALLUCINOGENS	7.4%	6.6%	8.1%	0.36%	
UPPERS	6.5%	5.6%	7.4%	0.43%	
DOWNERS	7.3%	6.3%	8.2%	0.47%	
ROHYPNOL	2.9%	2.5%	3.4%	0.23%	
STEROIDS	2.6%	2.2%	3.0%	0.21%	
ECSTASY	6.7%	6.1%	7.4%	0.33%	
HEROIN	1.6%	1.3%	1.9%	0.15%	

Lifetime Use of Selected Substances - Male Grade 10					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	20.4%	18.8%	22.0%	0.80%	
CIGARETTES	17.5%	16.0%	19.0%	0.75%	
SMOKELESS TOBACCO	7.4%	6.4%	8.3%	0.49%	
ALCOHOL	36.2%	34.7%	37.7%	0.75%	
INHALANTS	4.8%	3.6%	5.9%	0.58%	
ANY ILLICIT DRUG	17.2%	16.1%	18.3%	0.55%	
MARIJUANA	15.1%	14.0%	16.1%	0.52%	
MARIJUANA ONLY	6.2%	5.5%	6.7%	0.30%	
COCAINE OR CRACK	3.7%	3.1%	4.3%	0.30%	
COCAINE	3.4%	2.8%	4.0%	0.30%	
CRACK	0.6%	0.5%	0.7%	0.06%	
HALLUCINOGENS	2.5%	2.1%	3.0%	0.23%	
UPPERS	2.6%	2.1%	3.1%	0.25%	
DOWNERS	3.5%	3.0%	3.9%	0.23%	
ROHYPNOL	1.0%	0.8%	1.3%	0.12%	
STEROIDS	1.1%	0.9%	1.4%	0.13%	
ECSTASY	2.6%	2.2%	3.0%	0.21%	
HEROIN	0.8%	0.6%	1.0%	0.11%	

Lifetime Use of Selected Substances - Male Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	47.8%	45.2%	50.3%	1.27%	
CIGARETTES	44.4%	42.0%	46.7%	1.17%	
SMOKELESS TOBACCO	21.7%	19.4%	24.1%	1.16%	
ALCOHOL	73.1%	71.2%	75.0%	0.93%	
INHALANTS	14.6%	13.2%	15.9%	0.67%	
ANY ILLICIT DRUG	42.7%	41.0%	44.4%	0.85%	
MARIJUANA	41.8%	40.0%	43.5%	0.88%	
MARIJUANA ONLY	21.9%	20.5%	23.3%	0.69%	
COCAINE OR CRACK	12.4%	11.1%	13.8%	0.68%	

Lifetime Use of Selected Substances - Male Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	25.2%	23.3%	27.1%	0.97%	
CIGARETTES	22.0%	20.3%	23.7%	0.86%	
SMOKELESS TOBACCO	9.8%	8.7%	10.9%	0.54%	
ALCOHOL	41.2%	39.2%	43.2%	1.01%	
INHALANTS	4.5%	3.5%	5.4%	0.47%	
ANY ILLICIT DRUG	21.1%	19.7%	22.5%	0.72%	
MARIJUANA	19.0%	17.7%	20.3%	0.66%	
MARIJUANA ONLY	7.4%	6.5%	8.2%	0.43%	
COCAINE OR CRACK	4.9%	4.2%	5.6%	0.33%	

COCAINE	12.2%	10.8%	13.6%	0.69%
CRACK	2.6%	2.2%	3.0%	0.19%
HALLUCINOGENS	8.9%	7.5%	10.3%	0.72%
UPPERS	9.4%	7.8%	10.9%	0.77%
DOWNERS	10.3%	8.7%	11.8%	0.78%
ROHYPNOL	3.0%	2.4%	3.6%	0.31%
STEROIDS	2.6%	2.2%	2.9%	0.18%
ECSTASY	8.0%	6.8%	9.2%	0.59%
HEROIN	1.8%	1.4%	2.2%	0.22%

COCAINE	4.7%	4.0%	5.3%	0.34%
CRACK	0.8%	0.6%	1.0%	0.11%
HALLUCINOGENS	2.8%	2.2%	3.4%	0.30%
UPPERS	4.0%	3.3%	4.7%	0.37%
DOWNERS	4.6%	3.7%	5.5%	0.46%
ROHYPNOL	1.4%	1.0%	1.7%	0.18%
STEROIDS	1.3%	1.0%	1.5%	0.12%
ECSTASY	2.8%	2.2%	3.4%	0.28%
HEROIN	0.6%	0.3%	0.8%	0.11%

Lifetime Use of Selected Substances - Male Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	54.3%	51.1%	57.5%	1.60%	
CIGARETTES	51.1%	47.9%	54.3%	1.59%	
SMOKELESS TOBACCO	25.4%	22.3%	28.6%	1.59%	
ALCOHOL	76.6%	74.6%	78.7%	1.02%	
INHALANTS	15.2%	14.0%	16.3%	0.57%	
ANY ILLICIT DRUG	46.5%	44.7%	48.4%	0.92%	
MARIJUANA	45.4%	43.6%	47.2%	0.90%	
MARIJUANA ONLY	21.9%	20.5%	23.3%	0.70%	
COCAINE OR CRACK	14.8%	13.0%	16.6%	0.90%	
COCAINE	14.6%	12.8%	16.4%	0.89%	
CRACK	2.9%	2.4%	3.4%	0.24%	
HALLUCINOGENS	11.4%	9.8%	12.9%	0.76%	
UPPERS	10.7%	9.0%	12.4%	0.85%	
DOWNERS	11.8%	10.2%	13.5%	0.82%	
ROHYPNOL	2.7%	2.0%	3.3%	0.31%	
STEROIDS	3.1%	2.6%	3.7%	0.27%	
ECSTASY	9.6%	8.6%	10.7%	0.51%	
HEROIN	1.8%	1.4%	2.2%	0.19%	

Lifetime Use of Selected Substances - Male Grade 12					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	32.4%	29.6%	35.1%	1.38%	
CIGARETTES	29.4%	26.9%	31.8%	1.21%	
SMOKELESS TOBACCO	12.0%	9.9%	14.1%	1.06%	
ALCOHOL	48.8%	46.0%	51.6%	1.40%	
INHALANTS	3.3%	2.6%	4.0%	0.35%	
ANY ILLICIT DRUG	22.6%	21.2%	23.9%	0.67%	
MARIJUANA	20.5%	19.2%	21.8%	0.65%	
MARIJUANA ONLY	7.0%	6.2%	7.8%	0.40%	
COCAINE OR CRACK	5.5%	4.6%	6.3%	0.42%	
COCAINE	5.4%	4.6%	6.2%	0.40%	
CRACK	0.7%	0.4%	0.9%	0.11%	
HALLUCINOGENS	3.5%	3.0%	3.9%	0.25%	
UPPERS	4.4%	3.5%	5.2%	0.44%	
DOWNERS	5.1%	4.1%	6.1%	0.50%	
ROHYPNOL	1.0%	0.7%	1.2%	0.13%	
STEROIDS	1.3%	1.0%	1.7%	0.18%	
ECSTASY	3.7%	3.0%	4.3%	0.34%	
HEROIN	0.7%	0.5%	1.0%	0.12%	

Lifetime Use of Selected Substances - Female Overall					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	33.5%	31.9%	35.0%	0.77%	
CIGARETTES	32.9%	31.4%	34.5%	0.77%	
SMOKELESS TOBACCO	5.4%	5.1%	5.7%	0.17%	
ALCOHOL	67.5%	66.0%	69.0%	0.76%	
INHALANTS	16.5%	15.7%	17.2%	0.39%	
ANY ILLICIT DRUG	25.9%	24.6%	27.2%	0.63%	
MARIJUANA	24.0%	22.7%	25.3%	0.63%	
MARIJUANA ONLY	12.1%	11.3%	12.9%	0.41%	
COCAINE OR CRACK	7.6%	6.9%	8.4%	0.37%	
COCAINE	7.0%	6.2%	7.7%	0.38%	
CRACK	2.4%	2.2%	2.6%	0.10%	
HALLUCINOGENS	3.8%	3.3%	4.2%	0.21%	
UPPERS	6.0%	5.3%	6.8%	0.38%	
DOWNERS	6.1%	5.4%	6.8%	0.35%	
ROHYPNOL	2.7%	2.4%	3.1%	0.17%	
STEROIDS	0.9%	0.8%	1.0%	0.05%	
ECSTASY	5.2%	4.8%	5.6%	0.21%	
HEROIN	1.5%	1.3%	1.6%	0.07%	

Past 30 Day Use of Selected Substances - Female Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	13.2%	12.3%	14.1%	0.44%
CIGARETTES	12.9%	12.0%	13.8%	0.45%
SMOKELESS TOBACCO	1.7%	1.5%	1.8%	0.07%
ALCOHOL	31.5%	30.3%	32.8%	0.64%
INHALANTS	5.9%	5.2%	6.6%	0.33%
ANY ILLICIT DRUG	11.3%	10.5%	12.0%	0.38%
MARIJUANA	9.7%	8.9%	10.4%	0.36%
MARIJUANA ONLY	3.6%	3.2%	3.9%	0.20%
COCAINE OR CRACK	2.6%	2.3%	2.9%	0.14%
COCAINE	2.4%	2.1%	2.6%	0.14%
CRACK	0.7%	0.6%	0.8%	0.04%
HALLUCINOGENS	1.0%	0.9%	1.2%	0.07%
UPPERS	2.4%	2.1%	2.8%	0.17%
DOWNERS	2.4%	2.1%	2.7%	0.15%
ROHYPNOL	0.9%	0.8%	1.0%	0.06%
STEROIDS	0.3%	0.2%	0.3%	0.02%
ECSTASY	1.6%	1.4%	1.8%	0.11%
HEROIN	0.4%	0.4%	0.5%	0.03%

Lifetime Use of Selected Substances - Female Grade 7					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	16.9%	15.4%	18.4%	0.74%	
CIGARETTES	16.5%	15.1%	18.0%	0.74%	
SMOKELESS TOBACCO	3.0%	2.5%	3.5%	0.24%	
ALCOHOL	47.2%	44.9%	49.5%	1.14%	
INHALANTS	18.4%	17.1%	19.6%	0.62%	
ANY ILLICIT DRUG	9.5%	8.3%	10.8%	0.64%	
MARIJUANA	7.8%	6.6%	9.0%	0.59%	
MARIJUANA ONLY	4.2%	3.5%	4.9%	0.35%	
COCAINE OR CRACK	3.2%	2.6%	3.8%	0.29%	
COCAINE	2.3%	1.8%	2.8%	0.24%	
CRACK	1.9%	1.6%	2.3%	0.17%	
HALLUCINOGENS	1.0%	0.8%	1.2%	0.11%	
UPPERS	0.9%	0.7%	1.1%	0.10%	
DOWNERS	0.9%	0.7%	1.1%	0.10%	
ROHYPNOL	1.1%	0.8%	1.5%	0.18%	

Past 30 Day Use of Selected Substances - Female Grade 7				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	5.4%	4.6%	6.1%	0.37%
CIGARETTES	5.1%	4.4%	5.8%	0.35%
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.12%
ALCOHOL	16.3%	14.9%	17.7%	0.71%
INHALANTS	8.0%	7.1%	8.8%	0.43%
ANY ILLICIT DRUG	4.3%	3.4%	5.1%	0.40%
MARIJUANA	3.4%	2.7%	4.1%	0.36%
MARIJUANA ONLY	1.5%	1.0%	1.9%	0.22%
COCAINE OR CRACK	1.0%	0.7%	1.3%	0.14%
COCAINE	0.6%	0.4%	0.9%	0.13%
CRACK	0.6%	0.4%	0.7%	0.08%
HALLUCINOGENS	0.4%	0.3%	0.6%	0.07%
UPPERS	0.5%	0.3%	0.6%	0.08%
DOWNERS	0.4%	0.3%	0.6%	0.07%
ROHYPNOL	0.6%	0.3%	0.8%	0.11%

STEROIDS	0.9%	0.7%	1.0%	0.06%	
ECSTASY	1.6%	1.3%	1.9%	0.16%	
HEROIN	1.1%	0.9%	1.2%	0.09%	

STEROIDS	0.3%	0.2%	0.4%	0.05%
ECSTASY	0.4%	0.3%	0.5%	0.06%
HEROIN	0.4%	0.3%	0.5%	0.05%

Lifetime Use of S	elected Substance	s - Female	Grade 8	
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	26.6%	24.7%	28.4%	0.94%
CIGARETTES	25.7%	23.8%	27.6%	0.95%
SMOKELESS TOBACCO	4.7%	4.2%	5.3%	0.28%
ALCOHOL	61.3%	59.0%	63.6%	1.14%
INHALANTS	21.1%	19.6%	22.6%	0.74%
ANY ILLICIT DRUG	17.0%	15.4%	18.7%	0.84%
MARIJUANA	15.1%	13.5%	16.7%	0.79%
MARIJUANA ONLY	7.9%	6.9%	8.8%	0.48%
COCAINE OR CRACK	5.2%	4.4%	6.0%	0.40%
COCAINE	4.2%	3.4%	4.9%	0.36%
CRACK	2.7%	2.3%	3.2%	0.22%
HALLUCINOGENS	2.0%	1.6%	2.4%	0.18%
UPPERS	2.9%	2.4%	3.3%	0.23%
DOWNERS	2.6%	2.1%	3.0%	0.23%
ROHYPNOL	2.4%	1.9%	2.9%	0.26%
STEROIDS	1.1%	0.9%	1.4%	0.12%
ECSTASY	3.3%	2.8%	3.8%	0.24%
HEROIN	1.9%	1.6%	2.1%	0.12%

Past 30 Day Use of	Selected Substance	es - Femal	e Grade 8	
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	8.5%	7.6%	9.4%	0.44%
CIGARETTES	8.1%	7.2%	9.1%	0.47%
SMOKELESS TOBACCO	1.5%	1.2%	1.7%	0.13%
ALCOHOL	23.6%	21.8%	25.4%	0.90%
INHALANTS	8.0%	6.9%	9.1%	0.55%
ANY ILLICIT DRUG	7.5%	6.5%	8.5%	0.50%
MARIJUANA	6.5%	5.5%	7.5%	0.49%
MARIJUANA ONLY	2.7%	2.2%	3.2%	0.24%
COCAINE OR CRACK	1.7%	1.4%	2.0%	0.16%
COCAINE	1.4%	1.1%	1.7%	0.14%
CRACK	0.8%	0.6%	1.0%	0.12%
HALLUCINOGENS	0.5%	0.3%	0.7%	0.08%
UPPERS	1.1%	0.8%	1.4%	0.13%
DOWNERS	1.0%	0.7%	1.3%	0.14%
ROHYPNOL	0.9%	0.6%	1.1%	0.12%
STEROIDS	0.3%	0.2%	0.4%	0.05%
ECSTASY	1.1%	0.8%	1.4%	0.14%
HEROIN	0.5%	0.3%	0.6%	0.06%

Lifetime Use of Selected Substances - Female Grade 9					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	34.6%	32.7%	36.5%	0.94%	
CIGARETTES	34.1%	32.2%	35.9%	0.94%	
SMOKELESS TOBACCO	5.9%	5.2%	6.5%	0.32%	
ALCOHOL	69.9%	68.0%	71.8%	0.95%	
INHALANTS	18.9%	17.6%	20.2%	0.66%	
ANY ILLICIT DRUG	25.4%	23.9%	26.9%	0.76%	
MARIJUANA	23.1%	21.5%	24.6%	0.76%	
MARIJUANA ONLY	11.5%	10.2%	12.7%	0.62%	
COCAINE OR CRACK	7.4%	6.6%	8.3%	0.43%	
COCAINE	6.7%	5.8%	7.6%	0.43%	
CRACK	2.5%	2.1%	2.8%	0.18%	
HALLUCINOGENS	3.4%	3.0%	3.9%	0.23%	

Past 30 Day Use	Past 30 Day Use of Selected Substances - Female Grade 9					
CATEGORY	Prevalence	Boun	daries	Standard		
		Lower	Upper	Error		
TOBACCO	13.1%	12.0%	14.1%	0.53%		
CIGARETTES	12.6%	11.6%	13.7%	0.53%		
SMOKELESS TOBACCO	2.0%	1.6%	2.3%	0.16%		
ALCOHOL	32.5%	30.9%	34.1%	0.81%		
INHALANTS	6.9%	5.8%	7.9%	0.52%		
ANY ILLICIT DRUG	11.5%	10.5%	12.4%	0.47%		
MARIJUANA	9.8%	8.9%	10.7%	0.45%		
MARIJUANA ONLY	3.9%	3.3%	4.5%	0.32%		
COCAINE OR CRACK	2.7%	2.2%	3.2%	0.23%		
COCAINE	2.4%	1.9%	2.8%	0.24%		
CRACK	0.8%	0.6%	1.0%	0.09%		
HALLUCINOGENS	1.1%	0.9%	1.4%	0.13%		

_				
UPPERS	5.6%	4.9%	6.3%	0.36%
DOWNERS	5.4%	4.9%	6.0%	0.27%
ROHYPNOL	3.1%	2.7%	3.5%	0.21%
STEROIDS	0.9%	0.7%	1.2%	0.11%
ECSTASY	5.3%	4.8%	5.8%	0.25%
HEROIN	1.8%	1.4%	2.2%	0.20%

UPPERS	2.1%	1.7%	2.4%	0.19%
DOWNERS	2.1%	1.8%	2.5%	0.18%
ROHYPNOL	1.1%	0.8%	1.3%	0.11%
STEROIDS	0.3%	0.2%	0.3%	0.05%
ECSTASY	1.9%	1.5%	2.2%	0.17%
HEROIN	0.6%	0.3%	0.8%	0.11%

Lifetime Use of Selected Substances - Female Grade 10				
CATEGORY	Prevalence		daries	Standard
CATEGORY	Prevalence			
		Lower	Upper	Error
TOBACCO	38.8%	36.9%	40.7%	0.97%
CIGARETTES	38.3%	36.4%	40.2%	0.95%
SMOKELESS TOBACCO	6.2%	5.6%	6.8%	0.31%
ALCOHOL	74.8%	73.0%	76.5%	0.86%
INHALANTS	16.4%	15.0%	17.7%	0.67%
ANY ILLICIT DRUG	32.1%	30.2%	34.0%	0.93%
MARIJUANA	30.1%	28.2%	32.0%	0.94%
MARIJUANA ONLY	14.4%	13.1%	15.6%	0.62%
COCAINE OR CRACK	9.3%	8.1%	10.4%	0.59%
COCAINE	8.8%	7.6%	10.1%	0.60%
CRACK	2.6%	2.2%	3.0%	0.22%
HALLUCINOGENS	5.7%	4.9%	6.5%	0.40%
UPPERS	8.6%	7.4%	9.8%	0.59%
DOWNERS	8.6%	7.6%	9.7%	0.52%
ROHYPNOL	3.8%	3.1%	4.4%	0.33%
STEROIDS	1.0%	0.7%	1.2%	0.11%
ECSTASY	6.8%	6.0%	7.6%	0.39%
HEROIN	1.4%	1.1%	1.7%	0.14%

Past 30 Day Use of Selected Substances - Female Grade 10					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	15.4%	14.1%	16.7%	0.65%	
CIGARETTES	15.0%	13.8%	16.3%	0.64%	
SMOKELESS TOBACCO	1.9%	1.6%	2.2%	0.15%	
ALCOHOL	37.4%	35.7%	39.2%	0.88%	
INHALANTS	5.3%	4.3%	6.2%	0.46%	
ANY ILLICIT DRUG	14.3%	13.3%	15.4%	0.53%	
MARIJUANA	11.9%	10.8%	12.9%	0.53%	
MARIJUANA ONLY	3.8%	3.3%	4.3%	0.26%	
COCAINE OR CRACK	3.3%	2.6%	3.9%	0.32%	
COCAINE	3.1%	2.5%	3.7%	0.31%	
CRACK	0.9%	0.6%	1.1%	0.11%	
HALLUCINOGENS	1.6%	1.2%	2.0%	0.21%	
UPPERS	4.0%	3.4%	4.7%	0.32%	
DOWNERS	4.0%	3.4%	4.7%	0.34%	
ROHYPNOL	1.3%	1.0%	1.6%	0.15%	
STEROIDS	0.3%	0.2%	0.4%	0.05%	
ECSTASY	2.2%	1.7%	2.6%	0.24%	
HEROIN	0.4%	0.3%	0.5%	0.06%	

Lifetime Use of Selected Substances - Female Grade 11					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	42.1%	40.1%	44.1%	0.98%	
CIGARETTES	41.7%	39.7%	43.6%	0.98%	
SMOKELESS TOBACCO	6.2%	5.5%	6.9%	0.33%	
ALCOHOL	76.9%	75.4%	78.3%	0.73%	
INHALANTS	12.1%	11.0%	13.1%	0.51%	
ANY ILLICIT DRUG	37.1%	35.5%	38.6%	0.77%	
MARIJUANA	35.1%	33.4%	36.7%	0.81%	
MARIJUANA ONLY	17.4%	16.2%	18.5%	0.56%	
COCAINE OR CRACK	10.8%	9.3%	12.2%	0.72%	

Past 30 Day Use of Selected Substances - Female Grade 11					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	18.8%	17.3%	20.2%	0.72%	
CIGARETTES	18.4%	16.9%	19.8%	0.72%	
SMOKELESS TOBACCO	1.8%	1.5%	2.0%	0.12%	
ALCOHOL	39.5%	37.7%	41.2%	0.87%	
INHALANTS	3.3%	2.6%	3.9%	0.31%	
ANY ILLICIT DRUG	15.7%	14.3%	17.2%	0.73%	
MARIJUANA	13.7%	12.3%	15.1%	0.71%	
MARIJUANA ONLY	4.6%	3.9%	5.3%	0.36%	
COCAINE OR CRACK	3.8%	3.2%	4.4%	0.31%	

COCAINE	10.5%	9.1%	11.9%	0.70%
CRACK	2.4%	2.0%	2.9%	0.23%
HALLUCINOGENS	5.4%	4.5%	6.3%	0.44%
UPPERS	10.1%	8.7%	11.6%	0.73%
DOWNERS	10.5%	9.1%	11.8%	0.68%
ROHYPNOL	3.1%	2.6%	3.5%	0.23%
STEROIDS	0.6%	0.5%	0.8%	0.09%
ECSTASY	7.3%	6.1%	8.4%	0.56%
HEROIN	1.4%	1.1%	1.6%	0.13%

COCAINE	3.7%	3.1%	4.3%	0.30%
CRACK	0.5%	0.3%	0.7%	0.08%
HALLUCINOGENS	1.4%	1.0%	1.7%	0.20%
UPPERS	4.1%	3.4%	4.8%	0.37%
DOWNERS	4.1%	3.5%	4.8%	0.33%
ROHYPNOL	0.9%	0.7%	1.1%	0.09%
STEROIDS	0.2%	0.1%	0.2%	0.04%
ECSTASY	2.1%	1.6%	2.7%	0.26%
HEROIN	0.3%	0.1%	0.4%	0.06%

Lifetime Use of Selected Substances - Female Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	46.1%	43.6%	48.7%	1.27%	
CIGARETTES	45.7%	43.1%	48.3%	1.29%	
SMOKELESS TOBACCO	6.8%	5.9%	7.8%	0.48%	
ALCOHOL	79.1%	77.0%	81.1%	1.01%	
INHALANTS	9.4%	8.5%	10.2%	0.43%	
ANY ILLICIT DRUG	39.5%	37.4%	41.6%	1.04%	
MARIJUANA	38.3%	36.1%	40.4%	1.07%	
MARIJUANA ONLY	19.9%	18.6%	21.2%	0.65%	
COCAINE OR CRACK	11.3%	9.9%	12.7%	0.70%	
COCAINE	11.0%	9.6%	12.4%	0.71%	
CRACK	2.2%	1.7%	2.6%	0.21%	
HALLUCINOGENS	5.9%	5.1%	6.6%	0.37%	
UPPERS	9.8%	8.3%	11.4%	0.79%	
DOWNERS	10.1%	8.6%	11.6%	0.75%	
ROHYPNOL	3.1%	2.6%	3.5%	0.22%	
STEROIDS	0.7%	0.5%	0.9%	0.10%	
ECSTASY	8.2%	7.2%	9.1%	0.46%	
HEROIN	1.1%	0.8%	1.4%	0.13%	

Past 30 Day Use of Selected Substances - Female Grade 12					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	20.8%	18.9%	22.6%	0.93%	
CIGARETTES	20.6%	18.7%	22.4%	0.91%	
SMOKELESS TOBACCO	2.1%	1.6%	2.6%	0.25%	
ALCOHOL	44.2%	41.6%	46.7%	1.27%	
INHALANTS	2.7%	2.0%	3.4%	0.34%	
ANY ILLICIT DRUG	16.3%	14.9%	17.7%	0.69%	
MARIJUANA	14.5%	13.2%	15.7%	0.63%	
MARIJUANA ONLY	5.5%	4.9%	6.1%	0.30%	
COCAINE OR CRACK	3.6%	3.1%	4.1%	0.27%	
COCAINE	3.5%	2.9%	4.0%	0.27%	
CRACK	0.5%	0.3%	0.7%	0.09%	
HALLUCINOGENS	1.4%	1.2%	1.6%	0.12%	
UPPERS	3.4%	2.6%	4.2%	0.40%	
DOWNERS	3.6%	2.9%	4.2%	0.33%	
ROHYPNOL	0.9%	0.6%	1.1%	0.11%	
STEROIDS	0.3%	0.2%	0.4%	0.07%	
ECSTASY	2.2%	1.7%	2.7%	0.24%	
HEROIN	0.3%	0.2%	0.5%	0.07%	

Lifetime Use of Selected Substances - White Overall					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	36.0%	33.9%	38.1%	1.05%	
CIGARETTES	33.1%	31.2%	35.0%	0.97%	
SMOKELESS TOBACCO	13.8%	12.5%	15.1%	0.67%	
ALCOHOL	65.4%	63.5%	67.3%	0.94%	
INHALANTS	15.9%	15.1%	16.7%	0.41%	
ANY ILLICIT DRUG	26.6%	25.1%	28.1%	0.76%	
MARIJUANA	25.1%	23.7%	26.6%	0.72%	
MARIJUANA ONLY	12.2%	11.5%	12.9%	0.36%	
COCAINE OR CRACK	6.5%	5.9%	7.2%	0.34%	
COCAINE	6.0%	5.3%	6.7%	0.34%	
CRACK	2.0%	1.8%	2.2%	0.09%	
HALLUCINOGENS	6.5%	6.0%	6.9%	0.22%	
UPPERS	8.3%	7.5%	9.1%	0.40%	
DOWNERS	8.1%	7.2%	8.9%	0.41%	
ROHYPNOL	2.0%	1.8%	2.2%	0.11%	
STEROIDS	1.7%	1.5%	1.8%	0.08%	
ECSTASY	6.0%	5.4%	6.6%	0.31%	
HEROIN	1.4%	1.2%	1.6%	0.09%	

Past 30 Day Use of Selected Substances - White Overall					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.9%	16.5%	19.2%	0.67%	
CIGARETTES	15.7%	14.6%	16.9%	0.57%	
SMOKELESS TOBACCO	5.6%	4.9%	6.3%	0.35%	
ALCOHOL	32.3%	30.7%	33.9%	0.80%	
INHALANTS	4.6%	3.9%	5.4%	0.38%	
ANY ILLICIT DRUG	12.5%	11.6%	13.3%	0.41%	
MARIJUANA	10.9%	10.1%	11.7%	0.40%	
MARIJUANA ONLY	3.5%	3.2%	3.8%	0.16%	
COCAINE OR CRACK	2.2%	1.9%	2.4%	0.12%	
COCAINE	2.0%	1.8%	2.2%	0.11%	
CRACK	0.4%	0.4%	0.5%	0.04%	
HALLUCINOGENS	1.9%	1.7%	2.1%	0.10%	
UPPERS	3.4%	3.0%	3.8%	0.19%	
DOWNERS	3.4%	3.0%	3.8%	0.20%	
ROHYPNOL	0.8%	0.6%	0.9%	0.06%	
STEROIDS	0.6%	0.5%	0.7%	0.04%	
ECSTASY	2.1%	1.7%	2.4%	0.18%	
HEROIN	0.4%	0.3%	0.5%	0.05%	

Lifetime Use of Selected Substances - White Grade 7					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	15.1%	13.0%	17.2%	1.06%	
CIGARETTES	13.5%	11.5%	15.4%	0.98%	
SMOKELESS TOBACCO	4.8%	4.0%	5.7%	0.43%	
ALCOHOL	41.8%	39.3%	44.3%	1.25%	
INHALANTS	14.4%	12.4%	16.4%	0.98%	
ANY ILLICIT DRUG	6.6%	5.4%	7.8%	0.60%	
MARIJUANA	5.4%	4.4%	6.4%	0.50%	
MARIJUANA ONLY	3.0%	2.4%	3.7%	0.33%	
COCAINE OR CRACK	1.8%	1.3%	2.2%	0.21%	
COCAINE	1.1%	0.8%	1.5%	0.17%	
CRACK	1.3%	1.0%	1.6%	0.16%	
HALLUCINOGENS	1.1%	0.8%	1.4%	0.14%	
UPPERS	0.9%	0.6%	1.2%	0.15%	
DOWNERS	0.8%	0.5%	1.0%	0.10%	
ROHYPNOL	0.6%	0.4%	0.9%	0.11%	

Past 30 Day Use of Selected Substances - White Grade 7				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	4.6%	3.6%	5.5%	0.48%
CIGARETTES	3.7%	3.0%	4.5%	0.38%
SMOKELESS TOBACCO	1.6%	1.1%	2.0%	0.23%
ALCOHOL	11.7%	10.2%	13.2%	0.76%
INHALANTS	5.2%	4.2%	6.3%	0.54%
ANY ILLICIT DRUG	2.3%	1.7%	2.9%	0.31%
MARIJUANA	1.7%	1.2%	2.3%	0.26%
MARIJUANA ONLY	0.8%	0.5%	1.0%	0.14%
COCAINE OR CRACK	0.3%	0.2%	0.5%	0.07%
COCAINE	0.2%	0.1%	0.2%	0.04%
CRACK	0.2%	0.1%	0.3%	0.05%
HALLUCINOGENS	0.3%	0.1%	0.4%	0.07%
UPPERS	0.3%	0.2%	0.5%	0.08%
DOWNERS	0.3%	0.1%	0.4%	0.07%
ROHYPNOL	0.3%	0.2%	0.4%	0.06%

STEROIDS	1.4%	1.1%	1.7%	0.14%
ECSTASY	1.2%	0.9%	1.4%	0.12%
HEROIN	0.8%	0.6%	1.0%	0.10%

STEROIDS	0.4%	0.3%	0.6%	0.07%
ECSTASY	0.4%	0.3%	0.5%	0.06%
HEROIN	0.3%	0.2%	0.4%	0.06%

Lifetime Use of Selected Substances - White Grade 8				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	23.9%	21.6%	26.2%	1.14%
CIGARETTES	21.5%	19.3%	23.6%	1.10%
SMOKELESS TOBACCO	8.0%	6.7%	9.4%	0.67%
ALCOHOL	55.1%	52.0%	58.1%	1.53%
INHALANTS	18.2%	16.7%	19.7%	0.76%
ANY ILLICIT DRUG	13.3%	11.6%	15.0%	0.84%
MARIJUANA	12.0%	10.3%	13.7%	0.84%
MARIJUANA ONLY	6.2%	5.3%	7.0%	0.44%
COCAINE OR CRACK	2.8%	2.2%	3.3%	0.25%
COCAINE	2.1%	1.7%	2.5%	0.20%
CRACK	1.2%	0.9%	1.5%	0.16%
HALLUCINOGENS	3.1%	2.5%	3.6%	0.29%
UPPERS	2.9%	2.4%	3.5%	0.28%
DOWNERS	2.7%	2.2%	3.1%	0.24%
ROHYPNOL	1.5%	1.2%	1.8%	0.17%
STEROIDS	1.4%	0.8%	1.9%	0.27%
ECSTASY	2.8%	2.3%	3.2%	0.24%
HEROIN	1.3%	1.0%	1.6%	0.16%

Past 30 Day Use of Selected Substances - White Grade 8				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	8.6%	7.2%	9.9%	0.67%
CIGARETTES	7.2%	6.1%	8.3%	0.53%
SMOKELESS TOBACCO	2.7%	1.9%	3.6%	0.43%
ALCOHOL	19.1%	17.2%	21.0%	0.95%
INHALANTS	6.4%	5.1%	7.7%	0.64%
ANY ILLICIT DRUG	5.7%	4.7%	6.7%	0.50%
MARIJUANA	5.0%	4.0%	5.9%	0.49%
MARIJUANA ONLY	1.8%	1.3%	2.3%	0.24%
COCAINE OR CRACK	0.9%	0.7%	1.1%	0.12%
COCAINE	0.8%	0.6%	1.0%	0.11%
CRACK	0.2%	0.1%	0.3%	0.04%
HALLUCINOGENS	0.9%	0.6%	1.2%	0.14%
UPPERS	1.1%	0.8%	1.4%	0.14%
DOWNERS	1.1%	0.7%	1.4%	0.17%
ROHYPNOL	0.6%	0.5%	0.7%	0.07%
STEROIDS	0.4%	0.2%	0.6%	0.10%
ECSTASY	1.0%	0.7%	1.3%	0.15%
HEROIN	0.3%	0.1%	0.4%	0.07%

Lifetime Use of Selected Substances - White Grade 9				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	34.6%	31.4%	37.7%	1.56%
CIGARETTES	31.1%	28.1%	34.0%	1.45%
SMOKELESS TOBACCO	12.5%	10.5%	14.3%	0.95%
ALCOHOL	66.3%	64.1%	68.4%	1.08%
INHALANTS	17.3%	15.7%	19.0%	0.83%
ANY ILLICIT DRUG	23.6%	21.6%	25.7%	1.02%
MARIJUANA	21.6%	19.7%	23.6%	0.97%
MARIJUANA ONLY	11.3%	10.1%	12.6%	0.62%
COCAINE OR CRACK	4.7%	3.9%	5.6%	0.40%
COCAINE	4.0%	3.3%	4.8%	0.38%
CRACK	2.0%	1.5%	2.4%	0.22%
HALLUCINOGENS	5.3%	4.6%	6.1%	0.37%

Past 30 Day Use of Selected Substances - White Grade 9					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	15.7%	13.9%	17.5%	0.90%	
CIGARETTES	13.1%	11.6%	14.6%	0.74%	
SMOKELESS TOBACCO	5.0%	4.0%	6.1%	0.52%	
ALCOHOL	30.8%	28.7%	32.9%	1.05%	
INHALANTS	5.1%	4.1%	6.1%	0.50%	
ANY ILLICIT DRUG	10.8%	9.6%	11.9%	0.56%	
MARIJUANA	9.1%	7.9%	10.3%	0.62%	
MARIJUANA ONLY	3.4%	2.7%	4.0%	0.32%	
COCAINE OR CRACK	1.6%	1.2%	2.0%	0.20%	
COCAINE	1.4%	1.1%	1.7%	0.17%	
CRACK	0.5%	0.3%	0.7%	0.11%	
HALLUCINOGENS	1.8%	1.5%	2.1%	0.16%	

UPPERS	6.7%	5.7%	7.7%	0.50%
DOWNERS	6.4%	5.3%	7.4%	0.51%
ROHYPNOL	2.2%	1.7%	2.7%	0.24%
STEROIDS	1.5%	1.2%	1.8%	0.16%
ECSTASY	4.7%	4.1%	5.3%	0.29%
HEROIN	1.4%	1.1%	1.7%	0.16%

UPPERS	2.6%	2.0%	3.1%	0.27%
DOWNERS	2.7%	2.2%	3.3%	0.27%
ROHYPNOL	1.0%	0.7%	1.3%	0.15%
STEROIDS	0.5%	0.3%	0.6%	0.08%
ECSTASY	1.8%	1.5%	2.1%	0.15%
HEROIN	0.4%	0.3%	0.6%	0.09%

Lifetime Use of S	Lifetime Use of Selected Substances - White Grade 10					
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	40.9%	38.5%	43.3%	1.20%		
CIGARETTES	37.5%	35.1%	39.8%	1.18%		
SMOKELESS TOBACCO	16.1%	14.6%	17.6%	0.77%		
ALCOHOL	72.3%	70.7%	74.0%	0.83%		
INHALANTS	17.0%	15.5%	18.5%	0.73%		
ANY ILLICIT DRUG	32.2%	30.0%	34.4%	1.10%		
MARIJUANA	30.4%	28.4%	32.5%	1.02%		
MARIJUANA ONLY	14.4%	13.2%	15.7%	0.62%		
COCAINE OR CRACK	7.8%	6.8%	8.7%	0.48%		
COCAINE	7.3%	6.3%	8.3%	0.50%		
CRACK	2.3%	1.9%	2.6%	0.17%		
HALLUCINOGENS	8.3%	7.5%	9.0%	0.36%		
UPPERS	10.8%	9.5%	12.0%	0.64%		
DOWNERS	10.5%	9.1%	11.9%	0.69%		
ROHYPNOL	2.7%	2.1%	3.2%	0.27%		
STEROIDS	2.1%	1.8%	2.4%	0.17%		
ECSTASY	7.7%	6.7%	8.8%	0.53%		
HEROIN	1.4%	1.1%	1.7%	0.16%		

Past 30 Day Use of Selected Substances - White Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	21.0%	19.3%	22.7%	0.85%	
CIGARETTES	18.2%	16.6%	19.9%	0.83%	
SMOKELESS TOBACCO	6.6%	5.8%	7.4%	0.39%	
ALCOHOL	38.4%	36.7%	40.1%	0.84%	
INHALANTS	4.5%	3.3%	5.8%	0.62%	
ANY ILLICIT DRUG	15.6%	14.2%	17.0%	0.71%	
MARIJUANA	13.1%	11.8%	14.3%	0.63%	
MARIJUANA ONLY	4.1%	3.6%	4.7%	0.27%	
COCAINE OR CRACK	2.7%	2.1%	3.2%	0.27%	
COCAINE	2.5%	2.0%	3.0%	0.26%	
CRACK	0.6%	0.4%	0.8%	0.09%	
HALLUCINOGENS	2.4%	1.9%	2.9%	0.26%	
UPPERS	5.0%	4.2%	5.7%	0.38%	
DOWNERS	5.1%	4.2%	5.9%	0.41%	
ROHYPNOL	0.9%	0.6%	1.1%	0.13%	
STEROIDS	0.7%	0.6%	0.9%	0.09%	
ECSTASY	2.8%	2.2%	3.5%	0.33%	
HEROIN	0.6%	0.4%	0.8%	0.10%	

Lifetime Use of Selected Substances - White Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	48.0%	45.8%	50.2%	1.09%	
CIGARETTES	44.9%	42.9%	47.0%	1.01%	
SMOKELESS TOBACCO	19.4%	17.9%	21.0%	0.78%	
ALCOHOL	76.1%	74.1%	78.0%	0.96%	
INHALANTS	14.6%	13.4%	15.7%	0.58%	
ANY ILLICIT DRUG	40.1%	37.9%	42.3%	1.10%	
MARIJUANA	38.7%	36.5%	41.0%	1.14%	
MARIJUANA ONLY	17.8%	16.6%	19.1%	0.63%	
COCAINE OR CRACK	10.9%	9.8%	12.0%	0.56%	

Past 30 Day Use of Selected Substances - White Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	26.9%	25.0%	28.7%	0.91%	
CIGARETTES	24.0%	22.4%	25.6%	0.80%	
SMOKELESS TOBACCO	8.3%	7.4%	9.3%	0.47%	
ALCOHOL	43.6%	41.9%	45.4%	0.87%	
INHALANTS	3.7%	2.8%	4.6%	0.45%	
ANY ILLICIT DRUG	20.0%	18.5%	21.5%	0.76%	
MARIJUANA	17.9%	16.3%	19.4%	0.79%	
MARIJUANA ONLY	5.2%	4.4%	6.0%	0.40%	
COCAINE OR CRACK	3.7%	3.3%	4.1%	0.20%	

COCAINE	10.6%	9.5%	11.7%	0.56%	
CRACK	2.6%	2.3%	3.0%	0.19%	
HALLUCINOGENS	10.2%	9.0%	11.3%	0.59%	
UPPERS	14.3%	12.6%	15.9%	0.80%	
DOWNERS	13.8%	12.4%	15.2%	0.70%	
ROHYPNOL	2.6%	1.9%	3.2%	0.32%	
STEROIDS	1.9%	1.6%	2.1%	0.13%	
ECSTASY	9.6%	8.3%	10.8%	0.62%	
HEROIN	2.1%	1.6%	2.5%	0.21%	

COCAINE	3.6%	3.2%	4.0%	0.20%
CRACK	0.5%	0.4%	0.7%	0.09%
HALLUCINOGENS	3.0%	2.3%	3.7%	0.34%
UPPERS	6.1%	5.2%	6.9%	0.43%
DOWNERS	5.9%	5.1%	6.7%	0.40%
ROHYPNOL	1.0%	0.7%	1.4%	0.17%
STEROIDS	0.8%	0.6%	0.9%	0.07%
ECSTASY	3.1%	2.4%	3.9%	0.38%
HEROIN	0.4%	0.2%	0.6%	0.11%

Lifetime Use of Selected Substances - White Grade 12					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	53.5%	50.8%	56.2%	1.36%	
CIGARETTES	50.5%	47.6%	53.4%	1.47%	
SMOKELESS TOBACCO	22.2%	20.1%	24.3%	1.07%	
ALCOHOL	80.1%	78.5%	81.6%	0.78%	
INHALANTS	13.2%	12.1%	14.4%	0.58%	
ANY ILLICIT DRUG	44.1%	41.6%	46.7%	1.28%	
MARIJUANA	43.0%	40.4%	45.6%	1.29%	
MARIJUANA ONLY	20.6%	19.1%	22.1%	0.75%	
COCAINE OR CRACK	11.7%	10.2%	13.2%	0.74%	
COCAINE	11.3%	9.8%	12.8%	0.75%	
CRACK	2.5%	2.0%	3.0%	0.26%	
HALLUCINOGENS	11.1%	10.1%	12.1%	0.50%	
UPPERS	14.3%	12.5%	16.1%	0.89%	
DOWNERS	14.5%	12.8%	16.1%	0.84%	
ROHYPNOL	2.4%	1.9%	2.9%	0.24%	
STEROIDS	1.9%	1.5%	2.3%	0.21%	
ECSTASY	10.1%	8.8%	11.5%	0.68%	
HEROIN	1.6%	1.3%	2.0%	0.18%	

Past 30 Day Use of Selected Substances - White Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	31.2%	29.1%	33.3%	1.06%
CIGARETTES	28.7%	26.9%	30.6%	0.93%
SMOKELESS TOBACCO	9.8%	8.0%	11.5%	0.87%
ALCOHOL	50.1%	47.4%	52.7%	1.32%
INHALANTS	2.6%	1.9%	3.2%	0.32%
ANY ILLICIT DRUG	20.6%	18.8%	22.4%	0.88%
MARIJUANA	18.8%	17.1%	20.5%	0.85%
MARIJUANA ONLY	5.7%	4.9%	6.5%	0.38%
COCAINE OR CRACK	4.0%	3.3%	4.7%	0.35%
COCAINE	3.8%	3.1%	4.5%	0.35%
CRACK	0.6%	0.3%	0.8%	0.14%
HALLUCINOGENS	3.0%	2.5%	3.4%	0.23%
UPPERS	5.3%	4.3%	6.3%	0.50%
DOWNERS	5.6%	4.6%	6.5%	0.47%
ROHYPNOL	0.8%	0.6%	1.0%	0.11%
STEROIDS	0.8%	0.5%	1.0%	0.11%
ECSTASY	3.3%	2.3%	4.3%	0.49%
HEROIN	0.6%	0.4%	0.9%	0.12%

Lifetime Use of Selected Substances - African American Overall					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	24.5%	22.8%	26.1%	0.84%	
CIGARETTES	23.5%	21.9%	25.1%	0.82%	
SMOKELESS TOBACCO	4.8%	4.2%	5.4%	0.30%	
ALCOHOL	62.9%	61.0%	64.8%	0.96%	
INHALANTS	11.5%	10.3%	12.7%	0.59%	
ANY ILLICIT DRUG	26.1%	24.4%	27.8%	0.83%	
MARIJUANA	25.6%	23.9%	27.2%	0.84%	
MARIJUANA ONLY	20.1%	18.9%	21.3%	0.60%	
COCAINE OR CRACK	1.9%	1.4%	2.4%	0.24%	
COCAINE	1.6%	1.1%	2.1%	0.23%	
CRACK	0.9%	0.7%	1.2%	0.13%	
HALLUCINOGENS	1.2%	0.9%	1.5%	0.16%	
UPPERS	1.7%	1.3%	2.1%	0.21%	
DOWNERS	2.2%	1.7%	2.7%	0.25%	
ROHYPNOL	0.7%	0.5%	0.9%	0.09%	
STEROIDS	1.4%	1.1%	1.6%	0.14%	
ECSTASY	2.9%	2.5%	3.3%	0.21%	
HEROIN	0.8%	0.5%	1.0%	0.11%	

Past 30 Day Use of Selected Substances - African American Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	7.4%	6.7%	8.1%	0.35%
CIGARETTES	6.7%	6.0%	7.4%	0.34%
SMOKELESS TOBACCO	1.7%	1.5%	1.9%	0.09%
ALCOHOL	24.8%	22.9%	26.7%	0.97%
INHALANTS	4.3%	3.4%	5.2%	0.45%
ANY ILLICIT DRUG	10.7%	9.7%	11.7%	0.52%
MARIJUANA	10.0%	9.0%	11.0%	0.48%
MARIJUANA ONLY	7.1%	6.4%	7.7%	0.34%
COCAINE OR CRACK	0.8%	0.6%	1.1%	0.12%
COCAINE	0.6%	0.4%	0.9%	0.12%
CRACK	0.4%	0.3%	0.5%	0.07%
HALLUCINOGENS	0.5%	0.3%	0.6%	0.07%
UPPERS	0.7%	0.5%	0.9%	0.10%
DOWNERS	0.9%	0.7%	1.2%	0.12%
ROHYPNOL	0.4%	0.3%	0.4%	0.05%
STEROIDS	0.5%	0.4%	0.6%	0.06%
ECSTASY	1.1%	0.8%	1.4%	0.14%
HEROIN	0.4%	0.3%	0.5%	0.07%

Lifetime Use of Selected Substances - African American Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	18.3%	16.1%	20.5%	1.10%
CIGARETTES	17.7%	15.5%	19.8%	1.08%
SMOKELESS TOBACCO	3.3%	2.6%	4.1%	0.37%
ALCOHOL	49.7%	45.0%	54.3%	2.32%
INHALANTS	18.5%	16.1%	21.0%	1.22%
ANY ILLICIT DRUG	10.0%	8.0%	12.0%	1.00%
MARIJUANA	9.4%	7.4%	11.4%	1.00%
MARIJUANA ONLY	6.6%	4.9%	8.3%	0.84%
COCAINE OR CRACK	1.8%	1.3%	2.3%	0.25%
COCAINE	1.4%	0.9%	1.9%	0.26%
CRACK	1.0%	0.6%	1.3%	0.17%
HALLUCINOGENS	0.7%	0.4%	1.0%	0.14%
UPPERS	0.7%	0.3%	1.0%	0.18%
DOWNERS	0.5%	0.3%	0.6%	0.10%
ROHYPNOL	0.4%	0.2%	0.6%	0.11%

Past 30 Day Use of Selected Substances - African American Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	5.0%	4.4%	5.7%	0.33%
CIGARETTES	4.7%	4.0%	5.3%	0.33%
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.10%
ALCOHOL	16.5%	13.8%	19.1%	1.34%
INHALANTS	7.2%	5.8%	8.5%	0.68%
ANY ILLICIT DRUG	3.9%	3.0%	4.8%	0.46%
MARIJUANA	3.5%	2.7%	4.3%	0.42%
MARIJUANA ONLY	2.1%	1.5%	2.7%	0.31%
COCAINE OR CRACK	0.4%	0.2%	0.6%	0.10%
COCAINE	0.4%	0.2%	0.5%	0.10%
CRACK	0.3%	0.1%	0.4%	0.09%
HALLUCINOGENS	0.5%	0.3%	0.7%	0.11%
UPPERS	0.3%	0.1%	0.4%	0.09%
DOWNERS	0.2%	0.0%	0.4%	0.08%
ROHYPNOL	0.2%	0.0%	0.3%	0.07%

STEROIDS	1.6%	1.1%	2.0%	0.24%
ECSTASY	1.2%	0.8%	1.6%	0.20%
HEROIN	0.8%	0.4%	1.1%	0.18%

STEROIDS	0.5%	0.3%	0.7%	0.11%
ECSTASY	0.3%	0.1%	0.5%	0.10%
HEROIN	0.4%	0.2%	0.6%	0.11%

	10.1.			
Lifetime Use of Select	ed Substances - Af	rican Amer	ican Grade	8
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	22.2%	18.9%	25.6%	1.68%
CIGARETTES	20.8%	17.4%	24.1%	1.67%
SMOKELESS TOBACCO	4.8%	3.7%	5.9%	0.56%
ALCOHOL	61.4%	57.8%	65.1%	1.84%
INHALANTS	15.0%	13.4%	16.6%	0.81%
ANY ILLICIT DRUG	18.2%	16.1%	20.2%	1.03%
MARIJUANA	17.1%	14.9%	19.3%	1.11%
MARIJUANA ONLY	13.5%	11.9%	15.1%	0.82%
COCAINE OR CRACK	1.8%	1.0%	2.5%	0.36%
COCAINE	1.4%	0.8%	2.0%	0.28%
CRACK	1.1%	0.5%	1.7%	0.29%
HALLUCINOGENS	0.9%	0.6%	1.2%	0.15%
UPPERS	1.3%	0.8%	1.8%	0.26%
DOWNERS	1.3%	0.8%	1.8%	0.26%
ROHYPNOL	1.1%	0.7%	1.5%	0.18%
STEROIDS	1.5%	0.7%	2.3%	0.40%
ECSTASY	2.4%	1.5%	3.2%	0.43%
HEROIN	1.1%	0.7%	1.6%	0.23%

Past 30 Day Use of Sele	cted Substances - A	African Am	erican Grad	de 8
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	5.5%	4.4%	6.7%	0.60%
CIGARETTES	4.8%	3.5%	6.1%	0.66%
SMOKELESS TOBACCO	1.6%	1.4%	1.9%	0.14%
ALCOHOL	22.5%	19.7%	25.3%	1.40%
INHALANTS	5.2%	3.7%	6.7%	0.75%
ANY ILLICIT DRUG	7.1%	6.0%	8.2%	0.54%
MARIJUANA	6.4%	5.4%	7.4%	0.50%
MARIJUANA ONLY	4.4%	3.5%	5.4%	0.48%
COCAINE OR CRACK	0.8%	0.5%	1.1%	0.17%
COCAINE	0.6%	0.3%	0.9%	0.16%
CRACK	0.4%	0.1%	0.7%	0.13%
HALLUCINOGENS	0.3%	0.2%	0.5%	0.09%
UPPERS	0.6%	0.4%	0.8%	0.11%
DOWNERS	0.4%	0.2%	0.6%	0.08%
ROHYPNOL	0.5%	0.2%	0.8%	0.14%
STEROIDS	0.6%	0.2%	0.9%	0.18%
ECSTASY	1.0%	0.7%	1.4%	0.16%
HEROIN	0.6%	0.2%	0.9%	0.18%

Lifetime Use of Selected Substances - African American Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	27.3%	25.4%	29.2%	0.94%
CIGARETTES	26.1%	24.3%	27.9%	0.89%
SMOKELESS TOBACCO	5.3%	4.1%	6.5%	0.61%
ALCOHOL	64.4%	62.6%	66.1%	0.87%
INHALANTS	11.0%	8.4%	13.6%	1.29%
ANY ILLICIT DRUG	27.3%	25.1%	29.4%	1.07%
MARIJUANA	26.6%	24.5%	28.8%	1.09%
MARIJUANA ONLY	21.5%	19.9%	23.1%	0.80%
COCAINE OR CRACK	1.8%	1.1%	2.6%	0.39%
COCAINE	1.3%	0.6%	2.0%	0.35%
CRACK	1.0%	0.6%	1.5%	0.23%
HALLUCINOGENS	1.2%	0.6%	1.8%	0.29%

Post 20 Post the of Calenter I Ca				
Past 30 Day Use of Selected Substances - African American Grade 9				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	8.4%	7.0%	9.8%	0.71%
CIGARETTES	7.4%	5.9%	8.8%	0.73%
SMOKELESS TOBACCO	2.1%	1.5%	2.8%	0.33%
ALCOHOL	26.2%	22.6%	29.7%	1.78%
INHALANTS	4.2%	2.1%	6.4%	1.07%
ANY ILLICIT DRUG	10.9%	8.6%	13.1%	1.11%
MARIJUANA	9.7%	7.6%	11.9%	1.08%
MARIJUANA ONLY	7.2%	5.7%	8.6%	0.72%
COCAINE OR CRACK	1.2%	0.6%	1.7%	0.29%
COCAINE	0.6%	0.2%	1.1%	0.23%
CRACK	0.8%	0.3%	1.2%	0.22%
HALLUCINOGENS	0.5%	0.1%	0.9%	0.19%

•				
UPPERS	1.2%	0.6%	1.8%	0.29%
DOWNERS	1.5%	0.9%	2.1%	0.29%
ROHYPNOL	0.9%	0.6%	1.3%	0.18%
STEROIDS	0.7%	0.3%	1.1%	0.21%
ECSTASY	2.9%	1.8%	3.9%	0.51%
HEROIN	0.6%	0.3%	0.9%	0.17%

UPPERS	0.5%	0.3%	0.8%	0.11%
DOWNERS	0.7%	0.4%	1.1%	0.18%
ROHYPNOL	0.5%	0.2%	0.8%	0.16%
STEROIDS	0.2%	0.0%	0.4%	0.08%
ECSTASY	1.1%	0.3%	1.9%	0.41%
HEROIN	0.3%	0.0%	0.5%	0.13%

Lifetime Use of Selected Substances - African American Grade 10				
CATEGORY	Prevalence		daries	Standard
		Lower	Upper	Error
TOBACCO	26.2%	23.5%	29.0%	1.36%
CIGARETTES	25.7%	23.1%	28.2%	1.28%
SMOKELESS TOBACCO	4.9%	4.1%	5.6%	0.36%
ALCOHOL	68.1%	65.8%	70.3%	1.11%
INHALANTS	9.1%	7.8%	10.4%	0.67%
ANY ILLICIT DRUG	33.8%	31.9%	35.7%	0.95%
MARIJUANA	33.2%	31.5%	34.8%	0.83%
MARIJUANA ONLY	26.8%	24.9%	28.7%	0.97%
COCAINE OR CRACK	1.6%	0.9%	2.4%	0.37%
COCAINE	1.5%	0.8%	2.2%	0.35%
CRACK	0.4%	0.1%	0.7%	0.15%
HALLUCINOGENS	1.6%	0.9%	2.2%	0.34%
UPPERS	2.2%	1.2%	3.0%	0.45%
DOWNERS	2.9%	1.8%	4.0%	0.54%
ROHYPNOL	0.4%	0.1%	0.7%	0.15%
STEROIDS	1.4%	0.9%	1.8%	0.23%
ECSTASY	3.4%	2.7%	4.1%	0.35%
HEROIN	0.5%	0.2%	0.9%	0.18%

Past 30 Day Use of Selected Substances - African American Grade 10				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	7.3%	6.3%	8.3%	0.50%
CIGARETTES	6.8%	5.7%	7.9%	0.54%
SMOKELESS TOBACCO	2.0%	1.6%	2.4%	0.21%
ALCOHOL	27.2%	24.6%	29.7%	1.26%
INHALANTS	3.9%	2.7%	5.0%	0.58%
ANY ILLICIT DRUG	14.4%	12.9%	15.8%	0.74%
MARIJUANA	13.7%	12.3%	15.0%	0.67%
MARIJUANA ONLY	9.9%	8.7%	11.1%	0.60%
COCAINE OR CRACK	0.5%	0.2%	0.9%	0.18%
COCAINE	0.4%	0.1%	0.7%	0.15%
CRACK	0.2%	0.0%	0.4%	0.11%
HALLUCINOGENS	0.5%	0.1%	0.8%	0.18%
UPPERS	0.7%	0.3%	1.1%	0.21%
DOWNERS	1.5%	0.8%	2.2%	0.34%
ROHYPNOL	0.1%	0.0%	0.2%	0.05%
STEROIDS	0.5%	0.2%	0.7%	0.12%
ECSTASY	1.2%	0.8%	1.6%	0.21%
HEROIN	0.2%	0.0%	0.5%	0.12%

Lifetime Use of Selected Substances - African American Grade 11					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	26.1%	22.8%	29.4%	1.65%	
CIGARETTES	25.1%	22.0%	28.2%	1.54%	
SMOKELESS TOBACCO	5.3%	4.0%	6.7%	0.67%	
ALCOHOL	68.6%	65.7%	71.6%	1.48%	
INHALANTS	5.9%	4.3%	7.5%	0.80%	
ANY ILLICIT DRUG	37.2%	34.4%	40.0%	1.42%	
MARIJUANA	37.0%	34.1%	39.8%	1.41%	
MARIJUANA ONLY	29.5%	27.5%	31.5%	1.01%	
COCAINE OR CRACK	1.9%	0.9%	2.9%	0.49%	

Past 30 Day Use of Selected Substances - African American Grade 11						
CATEGORY	Prevalence	Boun	daries	Standard		
		Lower	Upper	Error		
TOBACCO	8.4%	6.8%	10.1%	0.82%		
CIGARETTES	7.9%	6.4%	9.5%	0.77%		
SMOKELESS TOBACCO	1.2%	0.7%	1.7%	0.26%		
ALCOHOL	27.7%	24.8%	30.6%	1.46%		
INHALANTS	2.2%	1.4%	3.0%	0.39%		
ANY ILLICIT DRUG	15.3%	13.1%	17.5%	1.10%		
MARIJUANA	14.5%	12.4%	16.5%	1.02%		
MARIJUANA ONLY	10.8%	9.2%	12.3%	0.77%		
COCAINE OR CRACK	1.0%	0.4%	1.6%	0.31%		

COCAINE	1.8%	0.9%	2.8%	0.49%
CRACK	0.6%	0.1%	1.1%	0.26%
HALLUCINOGENS	1.4%	0.5%	2.3%	0.46%
UPPERS	2.6%	1.6%	3.5%	0.50%
DOWNERS	3.4%	2.4%	4.4%	0.50%
ROHYPNOL	0.7%	0.2%	1.2%	0.24%
STEROIDS	1.6%	1.0%	2.3%	0.33%
ECSTASY	4.0%	2.8%	5.3%	0.62%
HEROIN	0.5%	0.0%	1.0%	0.23%

COCAINE	0.9%	0.3%	1.5%	0.30%
CRACK	0.4%	0.0%	0.7%	0.17%
HALLUCINOGENS	0.3%	0.0%	0.5%	0.11%
UPPERS	0.9%	0.4%	1.5%	0.27%
DOWNERS	1.5%	0.8%	2.2%	0.35%
ROHYPNOL	0.4%	0.1%	0.7%	0.15%
STEROIDS	0.9%	0.4%	1.3%	0.22%
ECSTASY	1.1%	0.6%	1.7%	0.28%
HEROIN	0.3%	0.0%	0.7%	0.18%

Lifetime Use of Selected Substances - African American Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	29.0%	25.6%	32.3%	1.69%	
CIGARETTES	28.2%	24.9%	31.5%	1.65%	
SMOKELESS TOBACCO	5.8%	4.1%	7.4%	0.82%	
ALCOHOL	69.7%	66.1%	73.2%	1.78%	
INHALANTS	5.5%	3.9%	7.1%	0.79%	
ANY ILLICIT DRUG	38.9%	36.1%	41.7%	1.40%	
MARIJUANA	38.9%	36.1%	41.6%	1.38%	
MARIJUANA ONLY	29.4%	27.3%	31.5%	1.04%	
COCAINE OR CRACK	2.7%	1.5%	3.9%	0.61%	
COCAINE	2.6%	1.4%	3.7%	0.59%	
CRACK	1.4%	0.5%	2.3%	0.46%	
HALLUCINOGENS	1.9%	1.1%	2.7%	0.41%	
UPPERS	3.1%	1.8%	4.3%	0.61%	
DOWNERS	4.9%	3.5%	6.3%	0.69%	
ROHYPNOL	0.7%	0.3%	1.0%	0.17%	
STEROIDS	1.5%	0.7%	2.3%	0.41%	
ECSTASY	4.6%	3.3%	5.9%	0.67%	
HEROIN	1.0%	0.3%	1.7%	0.35%	

Past 30 Day Use of Selected Substances - African American Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	11.2%	9.1%	13.2%	1.02%
CIGARETTES	10.4%	8.3%	12.5%	1.05%
SMOKELESS TOBACCO	2.4%	1.6%	3.2%	0.40%
ALCOHOL	32.7%	29.9%	35.4%	1.37%
INHALANTS	1.4%	0.7%	2.1%	0.36%
ANY ILLICIT DRUG	16.7%	13.8%	19.7%	1.48%
MARIJUANA	16.3%	13.4%	19.3%	1.49%
MARIJUANA ONLY	10.9%	9.1%	12.7%	0.90%
COCAINE OR CRACK	1.1%	0.5%	1.7%	0.31%
COCAINE	1.1%	0.5%	1.7%	0.31%
CRACK	0.4%	0.1%	0.6%	0.14%
HALLUCINOGENS	0.7%	0.3%	1.1%	0.20%
UPPERS	1.3%	0.5%	2.0%	0.37%
DOWNERS	1.6%	0.9%	2.3%	0.35%
ROHYPNOL	0.3%	0.1%	0.6%	0.13%
STEROIDS	0.6%	0.1%	1.1%	0.24%
ECSTASY	2.4%	1.3%	3.4%	0.52%
HEROIN	0.6%	0.1%	1.1%	0.27%

Lifetime Use of Selected Substances - Mexican American Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	40.0%	38.7%	41.3%	0.64%
CIGARETTES	39.2%	38.0%	40.4%	0.62%
SMOKELESS TOBACCO	8.3%	7.8%	8.8%	0.25%
ALCOHOL	69.2%	67.7%	70.7%	0.75%
INHALANTS	19.3%	18.2%	20.3%	0.52%
ANY ILLICIT DRUG	31.5%	29.7%	33.3%	0.89%
MARIJUANA	29.5%	27.6%	31.4%	0.94%
MARIJUANA ONLY	14.1%	12.8%	15.3%	0.62%
COCAINE OR CRACK	11.9%	11.2%	12.6%	0.34%
COCAINE	11.2%	10.5%	11.9%	0.36%
CRACK	3.3%	3.0%	3.5%	0.10%
HALLUCINOGENS	4.2%	3.6%	4.7%	0.28%
UPPERS	4.6%	3.9%	5.2%	0.32%
DOWNERS	5.3%	4.6%	6.0%	0.36%
ROHYPNOL	4.1%	3.6%	4.6%	0.25%
STEROIDS	1.3%	1.2%	1.5%	0.07%
ECSTASY	5.3%	4.9%	5.7%	0.20%
HEROIN	1.8%	1.6%	2.0%	0.09%

Past 30 Day Use of Selected Substances - Mexican American Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	16.3%	15.5%	17.1%	0.40%
CIGARETTES	15.6%	14.8%	16.3%	0.39%
SMOKELESS TOBACCO	3.2%	3.0%	3.4%	0.10%
ALCOHOL	34.7%	33.4%	35.9%	0.61%
INHALANTS	7.3%	6.5%	8.1%	0.40%
ANY ILLICIT DRUG	14.3%	13.3%	15.3%	0.49%
MARIJUANA	12.2%	11.2%	13.2%	0.51%
MARIJUANA ONLY	4.5%	4.0%	5.0%	0.25%
COCAINE OR CRACK	4.5%	4.1%	4.8%	0.18%
COCAINE	4.1%	3.8%	4.5%	0.17%
CRACK	1.0%	0.9%	1.1%	0.05%
HALLUCINOGENS	1.3%	1.1%	1.4%	0.08%
UPPERS	1.9%	1.6%	2.2%	0.15%
DOWNERS	2.2%	1.9%	2.6%	0.17%
ROHYPNOL	1.5%	1.3%	1.6%	0.09%
STEROIDS	0.5%	0.5%	0.6%	0.04%
ECSTASY	1.8%	1.6%	2.0%	0.09%
HEROIN	0.6%	0.5%	0.7%	0.04%

Lifetime Use of Selected Substances - Mexican American Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	22.3%	20.1%	24.4%	1.08%
CIGARETTES	21.7%	19.5%	23.8%	1.06%
SMOKELESS TOBACCO	4.6%	4.0%	5.3%	0.34%
ALCOHOL	51.3%	49.0%	53.6%	1.16%
INHALANTS	21.3%	19.9%	22.6%	0.68%
ANY ILLICIT DRUG	15.0%	12.8%	17.2%	1.11%
MARIJUANA	13.0%	10.9%	15.1%	1.04%
MARIJUANA ONLY	7.4%	6.2%	8.5%	0.56%
COCAINE OR CRACK	4.7%	3.8%	5.7%	0.47%
COCAINE	3.7%	2.9%	4.4%	0.37%
CRACK	2.6%	2.0%	3.2%	0.28%
HALLUCINOGENS	1.1%	0.8%	1.5%	0.19%
UPPERS	1.1%	0.8%	1.3%	0.13%
DOWNERS	1.2%	0.9%	1.5%	0.14%
ROHYPNOL	2.0%	1.5%	2.6%	0.27%

Past 30 Day Use of Selected Substances - Mexican American Grade 7					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	7.7%	6.6%	8.8%	0.54%	
CIGARETTES	7.2%	6.2%	8.2%	0.50%	
SMOKELESS TOBACCO	1.7%	1.4%	2.1%	0.19%	
ALCOHOL	20.2%	18.6%	21.8%	0.78%	
INHALANTS	9.6%	8.5%	10.6%	0.53%	
ANY ILLICIT DRUG	7.2%	5.8%	8.6%	0.72%	
MARIJUANA	5.9%	4.7%	7.1%	0.61%	
MARIJUANA ONLY	2.7%	2.1%	3.3%	0.30%	
COCAINE OR CRACK	1.8%	1.2%	2.3%	0.26%	
COCAINE	1.3%	0.9%	1.7%	0.22%	
CRACK	1.0%	0.7%	1.2%	0.14%	
HALLUCINOGENS	0.6%	0.3%	0.8%	0.12%	
UPPERS	0.6%	0.4%	0.8%	0.10%	
DOWNERS	0.6%	0.3%	0.8%	0.11%	
ROHYPNOL	0.9%	0.6%	1.2%	0.15%	

STEROIDS	1.1%	0.9%	1.3%	0.11%	
ECSTASY	1.8%	1.4%	2.1%	0.16%	
HEROIN	1.5%	1.2%	1.7%	0.11%	

STEROIDS	0.5%	0.3%	0.6%	0.07%
ECSTASY	0.6%	0.4%	0.8%	0.10%
HEROIN	0.7%	0.5%	0.9%	0.09%

Lifetime Use of Selected Substances - Mexican American Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	33.8%	31.6%	36.0%	1.09%	
CIGARETTES	32.9%	30.7%	35.1%	1.10%	
SMOKELESS TOBACCO	6.9%	6.1%	7.8%	0.43%	
ALCOHOL	65.3%	63.6%	67.1%	0.88%	
INHALANTS	23.7%	22.2%	25.2%	0.75%	
ANY ILLICIT DRUG	24.6%	22.2%	26.9%	1.16%	
MARIJUANA	22.6%	20.4%	24.8%	1.11%	
MARIJUANA ONLY	12.1%	10.5%	13.7%	0.81%	
COCAINE OR CRACK	8.2%	7.4%	9.0%	0.40%	
COCAINE	6.9%	6.1%	7.7%	0.39%	
CRACK	3.8%	3.3%	4.3%	0.26%	
HALLUCINOGENS	2.5%	2.0%	3.0%	0.24%	
UPPERS	2.4%	1.9%	2.9%	0.24%	
DOWNERS	2.5%	2.0%	3.1%	0.27%	
ROHYPNOL	3.7%	2.9%	4.5%	0.41%	
STEROIDS	1.6%	1.3%	2.0%	0.17%	
ECSTASY	3.5%	3.0%	4.1%	0.27%	
HEROIN	2.2%	1.8%	2.6%	0.18%	

Past 30 Day Use of Selected Substances - Mexican American Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	11.5%	10.3%	12.7%	0.60%	
CIGARETTES	11.1%	9.9%	12.2%	0.57%	
SMOKELESS TOBACCO	2.3%	1.8%	2.7%	0.21%	
ALCOHOL	27.7%	26.1%	29.2%	0.78%	
INHALANTS	9.3%	8.1%	10.6%	0.63%	
ANY ILLICIT DRUG	10.7%	9.5%	11.9%	0.61%	
MARIJUANA	9.3%	8.2%	10.5%	0.58%	
MARIJUANA ONLY	4.2%	3.6%	4.8%	0.32%	
COCAINE OR CRACK	2.8%	2.4%	3.2%	0.20%	
COCAINE	2.3%	2.0%	2.7%	0.19%	
CRACK	1.2%	1.0%	1.5%	0.13%	
HALLUCINOGENS	0.6%	0.4%	0.7%	0.09%	
UPPERS	0.9%	0.6%	1.1%	0.11%	
DOWNERS	0.9%	0.7%	1.0%	0.09%	
ROHYPNOL	1.3%	1.0%	1.6%	0.16%	
STEROIDS	0.6%	0.5%	0.8%	0.08%	
ECSTASY	1.1%	0.8%	1.4%	0.14%	
HEROIN	0.6%	0.4%	0.8%	0.10%	

Lifetime Use of Selected Substances - Mexican American Grade 9					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	42.0%	40.9%	43.2%	0.57%	
CIGARETTES	41.3%	40.1%	42.5%	0.60%	
SMOKELESS TOBACCO	9.2%	8.6%	9.8%	0.29%	
ALCOHOL	72.3%	70.5%	74.1%	0.89%	
INHALANTS	21.1%	19.4%	22.7%	0.81%	
ANY ILLICIT DRUG	34.0%	32.2%	35.8%	0.89%	
MARIJUANA	31.8%	29.8%	33.8%	1.00%	
MARIJUANA ONLY	15.2%	13.6%	16.8%	0.80%	
COCAINE OR CRACK	12.2%	11.2%	13.1%	0.45%	
COCAINE	11.6%	10.7%	12.5%	0.46%	
CRACK	3.3%	2.9%	3.7%	0.18%	
HALLUCINOGENS	4.3%	3.6%	4.9%	0.34%	

Past 30 Day Use of Selection	cted Substances - M	/lexican Am	nerican Gra	de 9
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	16.1%	15.2%	17.0%	0.43%
CIGARETTES	15.3%	14.5%	16.2%	0.41%
SMOKELESS TOBACCO	3.5%	3.1%	3.8%	0.17%
ALCOHOL	35.9%	34.2%	37.6%	0.85%
INHALANTS	7.9%	6.6%	9.2%	0.65%
ANY ILLICIT DRUG	15.8%	14.8%	16.7%	0.50%
MARIJUANA	13.7%	12.7%	14.7%	0.50%
MARIJUANA ONLY	5.3%	4.6%	6.0%	0.36%
COCAINE OR CRACK	4.8%	4.3%	5.4%	0.27%
COCAINE	4.5%	3.9%	5.0%	0.25%
CRACK	1.1%	0.9%	1.3%	0.12%
HALLUCINOGENS	1.2%	0.9%	1.5%	0.14%

UPPERS	4.7%	4.2%	5.1%	0.24%
DOWNERS	5.4%	4.9%	5.9%	0.24%
ROHYPNOL	4.7%	4.0%	5.3%	0.34%
STEROIDS	1.2%	1.0%	1.4%	0.11%
ECSTASY	5.9%	5.3%	6.5%	0.30%
HEROIN	2.3%	1.8%	2.9%	0.26%

UPPERS	1.8%	1.4%	2.1%	0.16%
DOWNERS	2.1%	1.8%	2.4%	0.16%
ROHYPNOL	1.7%	1.3%	2.0%	0.18%
STEROIDS	0.4%	0.3%	0.5%	0.06%
ECSTASY	2.2%	1.9%	2.5%	0.17%
HEROIN	0.7%	0.5%	1.0%	0.12%

Lifetime Use of Selected Substances - Mexican American Grade 10					
CATEGORY	Prevalence		daries	Standard	
CATEGORY	revalence	Lower	Upper	Error	
TOBACCO	47.1%	45.3%	48.9%	0.90%	
CIGARETTES	45.9%	44.1%	47.7%	0.90%	
SMOKELESS TOBACCO	9.2%	8.1%	10.3%	0.54%	
ALCOHOL	76.4%	74.7%	78.1%	0.87%	
INHALANTS	17.7%	16.1%	19.4%	0.82%	
ANY ILLICIT DRUG	38.3%	35.7%	40.9%	1.30%	
MARIJUANA	36.2%	33.4%	39.1%	1.44%	
MARIJUANA ONLY	15.9%	14.5%	17.3%	0.69%	
COCAINE OR CRACK	14.9%	13.8%	16.0%	0.55%	
COCAINE	14.3%	13.2%	15.5%	0.57%	
CRACK	3.6%	3.1%	4.1%	0.26%	
HALLUCINOGENS	6.4%	5.3%	7.6%	0.57%	
UPPERS	6.6%	5.4%	7.8%	0.61%	
DOWNERS	7.5%	6.4%	8.6%	0.57%	
ROHYPNOL	5.5%	4.6%	6.4%	0.46%	
STEROIDS	1.5%	1.1%	1.8%	0.17%	
ECSTASY	7.3%	6.5%	8.1%	0.40%	
HEROIN	1.9%	1.5%	2.2%	0.17%	

Past 30 Day Use of Selected Substances - Mexican American Grade 10					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	19.2%	18.1%	20.3%	0.56%	
CIGARETTES	18.4%	17.2%	19.5%	0.56%	
SMOKELESS TOBACCO	3.2%	2.7%	3.7%	0.24%	
ALCOHOL	40.8%	39.0%	42.6%	0.89%	
INHALANTS	6.0%	4.8%	7.3%	0.61%	
ANY ILLICIT DRUG	17.7%	16.1%	19.3%	0.79%	
MARIJUANA	15.0%	13.3%	16.7%	0.84%	
MARIJUANA ONLY	4.4%	3.8%	5.1%	0.33%	
COCAINE OR CRACK	5.8%	5.1%	6.4%	0.32%	
COCAINE	5.4%	4.8%	6.0%	0.30%	
CRACK	1.1%	0.9%	1.4%	0.13%	
HALLUCINOGENS	2.1%	1.7%	2.5%	0.20%	
UPPERS	2.8%	2.2%	3.4%	0.32%	
DOWNERS	3.4%	2.8%	4.0%	0.31%	
ROHYPNOL	1.9%	1.6%	2.3%	0.17%	
STEROIDS	0.7%	0.4%	1.0%	0.15%	
ECSTASY	2.5%	2.2%	2.9%	0.19%	
HEROIN	0.7%	0.5%	1.0%	0.11%	

Lifetime Use of Selected Substances - Mexican American Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	50.0%	48.0%	52.1%	1.02%	
CIGARETTES	49.2%	47.2%	51.2%	1.00%	
SMOKELESS TOBACCO	10.3%	9.2%	11.3%	0.52%	
ALCOHOL	78.2%	76.3%	80.1%	0.93%	
INHALANTS	14.2%	12.8%	15.6%	0.71%	
ANY ILLICIT DRUG	42.3%	40.1%	44.6%	1.13%	
MARIJUANA	40.2%	37.8%	42.6%	1.19%	
MARIJUANA ONLY	18.5%	16.6%	20.4%	0.96%	
COCAINE OR CRACK	17.3%	15.9%	18.6%	0.67%	

Past 30 Day Use of Selected Substances - Mexican American Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	22.6%	21.1%	24.1%	0.75%	
CIGARETTES	21.6%	20.1%	23.0%	0.72%	
SMOKELESS TOBACCO	4.4%	3.9%	4.8%	0.22%	
ALCOHOL	43.7%	41.7%	45.7%	0.99%	
INHALANTS	4.4%	3.2%	5.6%	0.59%	
ANY ILLICIT DRUG	18.3%	17.0%	19.7%	0.67%	
MARIJUANA	15.7%	14.2%	17.2%	0.74%	
MARIJUANA ONLY	5.1%	4.2%	6.0%	0.45%	
COCAINE OR CRACK	6.7%	5.9%	7.5%	0.38%	

COCAINE		17.0%	15.6%	18.3%	0.66%	
CRACK		3.2%	2.6%	3.8%	0.29%	
HALLUCINOGE	NS	5.5%	4.6%	6.3%	0.42%	
UPPERS		7.6%	6.3%	8.8%	0.62%	
DOWNERS		9.2%	7.5%	10.8%	0.82%	
ROHYPNOL		4.8%	4.0%	5.5%	0.36%	
STEROIDS		1.0%	0.8%	1.2%	0.11%	
ECSTASY		6.9%	6.0%	7.9%	0.48%	
HEROIN		1.5%	1.1%	1.8%	0.17%	

COCAINE	6.6%	5.8%	7.4%	0.38%
CRACK	0.8%	0.6%	1.0%	0.11%
HALLUCINOGENS	1.4%	1.1%	1.6%	0.13%
UPPERS	3.2%	2.6%	3.9%	0.33%
DOWNERS	4.1%	3.2%	5.0%	0.44%
ROHYPNOL	1.6%	1.3%	1.9%	0.16%
STEROIDS	0.4%	0.2%	0.5%	0.08%
ECSTASY	2.2%	1.7%	2.7%	0.24%
HEROIN	0.4%	0.2%	0.6%	0.08%

Lifetime Use of Selected Substances - Mexican American Grade 12				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	55.6%	52.9%	58.3%	1.36%
CIGARETTES	54.9%	52.3%	57.5%	1.29%
SMOKELESS TOBACCO	12.0%	10.6%	13.3%	0.68%
ALCOHOL	80.4%	78.0%	82.8%	1.20%
INHALANTS	13.2%	12.2%	14.2%	0.52%
ANY ILLICIT DRUG	44.8%	42.1%	47.6%	1.36%
MARIJUANA	43.2%	40.3%	46.1%	1.44%
MARIJUANA ONLY	19.1%	17.4%	20.9%	0.88%
COCAINE OR CRACK	19.4%	17.7%	21.0%	0.82%
COCAINE	19.3%	17.7%	21.0%	0.83%
CRACK	3.1%	2.5%	3.6%	0.27%
HALLUCINOGENS	7.4%	5.9%	8.9%	0.75%
UPPERS	7.9%	6.3%	9.5%	0.78%
DOWNERS	9.0%	7.1%	10.8%	0.93%
ROHYPNOL	4.6%	4.0%	5.3%	0.33%
STEROIDS	1.7%	1.2%	2.2%	0.27%
ECSTASY	8.5%	7.7%	9.3%	0.41%
HEROIN	1.2%	0.8%	1.7%	0.22%

Past 30 Day Use of Selected Substances - Mexican American Grade 12				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	27.6%	25.3%	30.0%	1.17%
CIGARETTES	26.8%	24.5%	29.1%	1.14%
SMOKELESS TOBACCO	5.0%	4.3%	5.8%	0.38%
ALCOHOL	49.6%	46.8%	52.4%	1.39%
INHALANTS	4.1%	3.1%	5.1%	0.48%
ANY ILLICIT DRUG	20.0%	18.3%	21.7%	0.85%
MARIJUANA	17.3%	15.7%	18.8%	0.79%
MARIJUANA ONLY	5.7%	5.1%	6.4%	0.33%
COCAINE OR CRACK	6.8%	6.0%	7.6%	0.38%
COCAINE	6.8%	6.0%	7.5%	0.38%
CRACK	0.6%	0.4%	0.7%	0.09%
HALLUCINOGENS	2.3%	1.8%	2.8%	0.25%
UPPERS	3.1%	2.3%	4.0%	0.43%
DOWNERS	3.8%	2.9%	4.7%	0.46%
ROHYPNOL	1.4%	1.1%	1.7%	0.15%
STEROIDS	0.8%	0.3%	1.2%	0.22%
ECSTASY	2.7%	2.2%	3.1%	0.23%
HEROIN	0.3%	0.1%	0.4%	0.08%

Lifetime Use of Selected Substances - Border Overall				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	38.0%	36.8%	39.3%	0.62%
CIGARETTES	37.4%	36.2%	38.5%	0.59%
SMOKELESS TOBACCO	7.6%	7.0%	8.3%	0.32%
ALCOHOL	67.3%	64.9%	69.7%	1.18%
INHALANTS	17.9%	16.7%	19.2%	0.62%
ANY ILLICIT DRUG	27.4%	25.9%	28.9%	0.76%
MARIJUANA	25.0%	23.4%	26.6%	0.80%
MARIJUANA ONLY	11.0%	9.9%	12.0%	0.51%
COCAINE OR CRACK	12.3%	11.5%	13.1%	0.40%
COCAINE	11.7%	10.9%	12.6%	0.41%
CRACK	3.3%	3.1%	3.5%	0.12%
HALLUCINOGENS	4.0%	3.1%	5.0%	0.47%
UPPERS	3.4%	3.1%	3.8%	0.18%
DOWNERS	3.8%	3.4%	4.2%	0.18%
ROHYPNOL	5.1%	4.1%	6.1%	0.50%
STEROIDS	1.2%	1.0%	1.5%	0.12%
ECSTASY	4.2%	3.9%	4.6%	0.18%
HEROIN	1.5%	1.4%	1.6%	0.05%

HEROIN	1.5%	1.4%	1.6%	0.05%
Lifetime Use of S	elected Substance	s - Border (Grade 7	
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	17.8%	15.4%	20.2%	1.19%
CIGARETTES	17.1%	15.0%	19.3%	1.06%
SMOKELESS TOBACCO	3.5%	3.0%	3.9%	0.23%
ALCOHOL	46.8%	43.6%	50.0%	1.61%
INHALANTS	18.0%	16.6%	19.4%	0.70%
ANY ILLICIT DRUG	9.9%	8.1%	11.7%	0.91%
MARIJUANA	8.2%	6.4%	9.9%	0.88%
MARIJUANA ONLY	4.7%	3.9%	5.5%	0.40%
COCAINE OR CRACK	3.2%	2.0%	4.4%	0.59%
COCAINE	2.6%	1.6%	3.7%	0.53%
CRACK	1.7%	1.0%	2.4%	0.36%
HALLUCINOGENS	0.9%	0.6%	1.1%	0.13%
UPPERS	0.9%	0.6%	1.2%	0.14%
DOWNERS	0.8%	0.6%	1.1%	0.13%
ROHYPNOL	1.9%	1.2%	2.6%	0.36%

Past 30-day Use of Selected Substances - Border Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	16.4%	0.5%	15.4%	17.41%
CIGARETTES	15.8%	0.5%	14.8%	16.76%
SMOKELESS TOBACCO	3.1%	0.2%	2.8%	3.37%
ALCOHOL	34.6%	0.8%	33.0%	36.21%
INHALANTS	7.4%	0.5%	6.3%	8.46%
ANY ILLICIT DRUG	12.3%	0.4%	11.4%	13.13%
MARIJUANA	10.0%	0.4%	9.2%	10.78%
MARIJUANA ONLY	3.3%	0.2%	2.9%	3.73%
COCAINE OR CRACK	4.9%	0.2%	4.5%	5.38%
COCAINE	4.6%	0.2%	4.2%	5.04%
CRACK	1.1%	0.1%	1.0%	1.19%
HALLUCINOGENS	1.3%	0.1%	1.0%	1.48%
UPPERS	1.3%	0.1%	1.1%	1.49%
DOWNERS	1.5%	0.1%	1.3%	1.69%
ROHYPNOL	1.8%	0.2%	1.4%	2.10%
STEROIDS	0.6%	0.1%	0.4%	0.68%
ECSTASY	1.5%	0.1%	1.3%	1.80%
HEROIN	0.5%	0.0%	0.5%	0.57%

Past 30-day Use of Selected Substances - Border Grade 7				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	6.0%	0.7%	4.6%	7.47%
CIGARETTES	5.5%	0.7%	4.2%	6.81%
SMOKELESS TOBACCO	1.4%	0.2%	1.1%	1.77%
ALCOHOL	17.4%	1.0%	15.4%	19.34%
INHALANTS	9.1%	0.7%	7.8%	10.42%
ANY ILLICIT DRUG	4.6%	0.5%	3.6%	5.65%
MARIJUANA	3.6%	0.5%	2.7%	4.56%
MARIJUANA ONLY	1.7%	0.2%	1.3%	2.10%
COCAINE OR CRACK	1.3%	0.4%	0.6%	1.99%
COCAINE	1.0%	0.3%	0.4%	1.67%
CRACK	0.7%	0.2%	0.3%	1.03%
HALLUCINOGENS	0.4%	0.1%	0.2%	0.59%
UPPERS	0.5%	0.1%	0.2%	0.69%
DOWNERS	0.4%	0.1%	0.2%	0.50%
ROHYPNOL	0.8%	0.2%	0.5%	1.17%

STEROIDS	1.0%	0.8%	1.2%	0.11%
ECSTASY	1.0%	0.7%	1.3%	0.15%
HEROIN	1.2%	0.8%	1.5%	0.19%

STEROIDS	0.5%	0.1%	0.3%	0.66%
ECSTASY	0.5%	0.1%	0.4%	0.62%
HEROIN	0.5%	0.1%	0.3%	0.72%

Lifetime Use of Selected Substances - Border Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	30.0%	27.9%	32.0%	1.02%
CIGARETTES	29.0%	26.9%	31.1%	1.06%
SMOKELESS TOBACCO	6.1%	5.0%	7.1%	0.52%
ALCOHOL	61.3%	59.6%	62.9%	0.84%
INHALANTS	20.8%	19.8%	21.9%	0.52%
ANY ILLICIT DRUG	19.3%	17.7%	21.0%	0.81%
MARIJUANA	17.5%	16.2%	18.9%	0.69%
MARIJUANA ONLY	8.1%	7.3%	8.9%	0.41%
COCAINE OR CRACK	8.3%	7.2%	9.3%	0.51%
COCAINE	7.3%	6.2%	8.4%	0.54%
CRACK	3.9%	3.4%	4.3%	0.24%
HALLUCINOGENS	2.2%	1.9%	2.6%	0.17%
UPPERS	1.9%	1.2%	2.5%	0.30%
DOWNERS	2.3%	1.4%	3.1%	0.42%
ROHYPNOL	4.3%	2.9%	5.7%	0.69%
STEROIDS	1.2%	0.8%	1.6%	0.20%
ECSTASY	2.6%	2.0%	3.2%	0.30%
HEROIN	1.7%	1.5%	1.9%	0.11%

Past 30-day Use of Selected Substances - Border Grade 8				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	10.3%	0.6%	9.1%	11.53%
CIGARETTES	9.8%	0.6%	8.5%	10.99%
SMOKELESS TOBACCO	2.3%	0.2%	1.9%	2.66%
ALCOHOL	25.7%	0.9%	23.9%	27.45%
INHALANTS	9.1%	0.9%	7.2%	10.97%
ANY ILLICIT DRUG	8.8%	0.7%	7.4%	10.20%
MARIJUANA	7.5%	0.6%	6.3%	8.67%
MARIJUANA ONLY	2.8%	0.2%	2.4%	3.26%
COCAINE OR CRACK	3.1%	0.3%	2.4%	3.70%
COCAINE	2.7%	0.3%	2.0%	3.26%
CRACK	1.3%	0.2%	0.8%	1.76%
HALLUCINOGENS	0.6%	0.1%	0.5%	0.75%
UPPERS	0.7%	0.1%	0.4%	0.96%
DOWNERS	0.8%	0.1%	0.5%	1.00%
ROHYPNOL	1.5%	0.3%	0.9%	1.95%
STEROIDS	0.6%	0.1%	0.4%	0.85%
ECSTASY	1.0%	0.2%	0.5%	1.50%
HEROIN	0.5%	0.1%	0.3%	0.65%

Lifetime Use of Selected Substances - Border Grade 9				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	40.0%	38.6%	41.4%	0.71%
CIGARETTES	39.7%	38.1%	41.2%	0.77%
SMOKELESS TOBACCO	8.6%	8.0%	9.1%	0.27%
ALCOHOL	70.6%	67.7%	73.5%	1.43%
INHALANTS	20.1%	17.4%	22.8%	1.34%
ANY ILLICIT DRUG	29.2%	27.1%	31.4%	1.08%
MARIJUANA	26.1%	24.2%	28.0%	0.95%
MARIJUANA ONLY	11.2%	10.1%	12.3%	0.55%
COCAINE OR CRACK	12.8%	11.5%	14.1%	0.66%
COCAINE	12.1%	10.9%	13.4%	0.62%
CRACK	3.4%	2.8%	4.0%	0.29%
HALLUCINOGENS	3.7%	2.7%	4.8%	0.52%

Past 30-day Use of Selected Substances - Border Grade 9				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	16.3%	0.7%	14.8%	17.69%
CIGARETTES	15.7%	0.7%	14.3%	17.00%
SMOKELESS TOBACCO	3.4%	0.2%	2.9%	3.81%
ALCOHOL	36.6%	1.0%	34.5%	38.58%
INHALANTS	7.5%	1.1%	5.3%	9.75%
ANY ILLICIT DRUG	13.2%	0.8%	11.6%	14.72%
MARIJUANA	10.9%	0.7%	9.5%	12.22%
MARIJUANA ONLY	3.5%	0.4%	2.8%	4.28%
COCAINE OR CRACK	5.3%	0.4%	4.5%	6.09%
COCAINE	4.8%	0.4%	4.1%	5.56%
CRACK	1.3%	0.2%	0.9%	1.65%
HALLUCINOGENS	1.3%	0.3%	0.8%	1.75%

_				
UPPERS	3.8%	3.5%	4.1%	0.17%
DOWNERS	4.4%	4.0%	4.8%	0.21%
ROHYPNOL	5.8%	4.5%	7.0%	0.63%
STEROIDS	1.0%	0.7%	1.3%	0.15%
ECSTASY	4.4%	3.8%	4.9%	0.26%
HEROIN	2.0%	1.2%	2.8%	0.41%

UPPERS	1.4%	0.2%	1.1%	1.77%
DOWNERS	1.7%	0.2%	1.4%	2.03%
ROHYPNOL	2.2%	0.3%	1.6%	2.83%
STEROIDS	0.3%	0.1%	0.1%	0.41%
ECSTASY	1.7%	0.2%	1.4%	2.01%
HEROIN	0.6%	0.1%	0.4%	0.81%

Lifetime Use of Se	elected Substances	- Border G	Frade 10	
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	44.1%	42.5%	45.8%	0.83%
CIGARETTES	43.3%	41.8%	44.8%	0.76%
SMOKELESS TOBACCO	8.1%	7.3%	8.8%	0.39%
ALCOHOL	74.4%	72.0%	76.7%	1.17%
INHALANTS	17.3%	15.2%	19.5%	1.08%
ANY ILLICIT DRUG	33.4%	30.5%	36.2%	1.41%
MARIJUANA	30.2%	27.3%	32.9%	1.40%
MARIJUANA ONLY	13.0%	11.1%	14.8%	0.94%
COCAINE OR CRACK	15.5%	14.8%	16.2%	0.35%
COCAINE	15.2%	14.4%	15.9%	0.36%
CRACK	4.0%	3.6%	4.4%	0.20%
HALLUCINOGENS	5.1%	3.7%	6.5%	0.70%
UPPERS	4.2%	3.5%	4.8%	0.35%
DOWNERS	4.8%	4.2%	5.4%	0.29%
ROHYPNOL	6.6%	4.8%	8.3%	0.86%
STEROIDS	1.4%	1.0%	1.7%	0.20%
ECSTASY	6.0%	5.3%	6.7%	0.35%
HEROIN	1.6%	1.3%	1.9%	0.13%

Past 30-day Use of Selected Substances - Border Grade 10					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	19.0%	0.6%	17.7%	20.22%	
CIGARETTES	18.3%	0.6%	17.1%	19.58%	
SMOKELESS TOBACCO	3.0%	0.2%	2.6%	3.42%	
ALCOHOL	40.1%	1.2%	37.6%	42.54%	
INHALANTS	6.1%	1.0%	4.1%	8.04%	
ANY ILLICIT DRUG	15.0%	0.8%	13.5%	16.61%	
MARIJUANA	11.8%	0.8%	10.2%	13.28%	
MARIJUANA ONLY	3.5%	0.4%	2.8%	4.18%	
COCAINE OR CRACK	6.9%	0.2%	6.4%	7.33%	
COCAINE	6.5%	0.2%	6.1%	6.93%	
CRACK	1.3%	0.2%	1.0%	1.65%	
HALLUCINOGENS	1.7%	0.2%	1.3%	2.05%	
UPPERS	1.5%	0.2%	1.1%	1.79%	
DOWNERS	2.1%	0.2%	1.7%	2.53%	
ROHYPNOL	2.3%	0.3%	1.7%	2.94%	
STEROIDS	0.8%	0.2%	0.4%	1.27%	
ECSTASY	2.0%	0.2%	1.6%	2.42%	
HEROIN	0.7%	0.1%	0.4%	0.93%	

Lifetime Use of Selected Substances - Border Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	49.7%	45.9%	53.4%	1.87%	
CIGARETTES	49.1%	45.5%	52.7%	1.80%	
SMOKELESS TOBACCO	9.0%	7.2%	10.7%	0.87%	
ALCOHOL	77.8%	74.2%	81.5%	1.82%	
INHALANTS	14.3%	12.0%	16.5%	1.13%	
ANY ILLICIT DRUG	38.2%	35.2%	41.3%	1.54%	
MARIJUANA	35.5%	32.3%	38.8%	1.62%	
MARIJUANA ONLY	15.3%	13.4%	17.1%	0.93%	
COCAINE OR CRACK	17.7%	16.3%	19.1%	0.71%	

Past 30-day Use of Selected Substances - Border Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	23.1%	0.8%	21.4%	24.70%	
CIGARETTES	22.3%	0.7%	20.9%	23.67%	
SMOKELESS TOBACCO	3.7%	0.3%	3.0%	4.30%	
ALCOHOL	45.1%	1.3%	42.5%	47.58%	
INHALANTS	5.6%	1.0%	3.6%	7.61%	
ANY ILLICIT DRUG	16.5%	0.6%	15.3%	17.73%	
MARIJUANA	13.3%	0.6%	12.2%	14.40%	
MARIJUANA ONLY	3.9%	0.3%	3.3%	4.50%	
COCAINE OR CRACK	7.0%	0.5%	5.9%	8.10%	

COCAINE	17.5%	16.0%	18.9%	0.73%
CRACK	3.4%	2.8%	4.0%	0.30%
HALLUCINOGENS	5.9%	4.0%	7.8%	0.94%
UPPERS	5.2%	4.0%	6.4%	0.60%
DOWNERS	5.6%	4.4%	6.8%	0.60%
ROHYPNOL	6.5%	5.2%	7.7%	0.62%
STEROIDS	1.3%	1.0%	1.5%	0.13%
ECSTASY	5.3%	4.2%	6.4%	0.55%
HEROIN	1.1%	0.7%	1.4%	0.17%

COCAINE	6.9%	0.6%	5.8%	8.05%
CRACK	1.0%	0.1%	0.7%	1.20%
HALLUCINOGENS	1.7%	0.2%	1.3%	2.16%
UPPERS	1.9%	0.3%	1.3%	2.50%
DOWNERS	2.0%	0.2%	1.6%	2.41%
ROHYPNOL	2.1%	0.2%	1.7%	2.48%
STEROIDS	0.7%	0.1%	0.4%	0.99%
ECSTASY	1.8%	0.4%	1.1%	2.52%
HEROIN	0.2%	0.1%	0.1%	0.35%

Lifetime Use of Se	Lifetime Use of Selected Substances - Border Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard		
		Lower	Upper	Error		
TOBACCO	55.7%	52.2%	59.1%	1.74%		
CIGARETTES	54.9%	51.8%	58.0%	1.54%		
SMOKELESS TOBACCO	12.4%	10.0%	14.8%	1.20%		
ALCOHOL	80.1%	75.3%	84.8%	2.38%		
INHALANTS	14.4%	12.7%	16.1%	0.83%		
ANY ILLICIT DRUG	42.3%	38.8%	45.9%	1.79%		
MARIJUANA	40.6%	36.6%	44.6%	2.01%		
MARIJUANA ONLY	16.9%	14.5%	19.3%	1.20%		
COCAINE OR CRACK	20.5%	18.3%	22.6%	1.08%		
COCAINE	20.4%	18.2%	22.5%	1.09%		
CRACK	3.7%	3.0%	4.4%	0.35%		
HALLUCINOGENS	8.4%	5.4%	11.3%	1.46%		
UPPERS	5.9%	4.6%	7.2%	0.64%		
DOWNERS	6.1%	4.6%	7.5%	0.73%		
ROHYPNOL	6.6%	5.3%	7.8%	0.62%		
STEROIDS	1.9%	1.3%	2.4%	0.27%		
ECSTASY	7.7%	6.7%	8.7%	0.50%		
HEROIN	1.3%	0.6%	2.0%	0.35%		

Past 30-day Use of Selected Substances - Border Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	30.3%	1.2%	27.9%	32.67%
CIGARETTES	29.6%	1.1%	27.3%	31.87%
SMOKELESS TOBACCO	5.6%	0.6%	4.4%	6.81%
ALCOHOL	51.5%	1.9%	47.6%	55.34%
INHALANTS	5.7%	0.7%	4.3%	7.11%
ANY ILLICIT DRUG	18.8%	0.9%	17.0%	20.63%
MARIJUANA	15.7%	0.8%	14.2%	17.24%
MARIJUANA ONLY	5.0%	0.4%	4.1%	5.88%
COCAINE OR CRACK	7.6%	0.5%	6.6%	8.61%
COCAINE	7.5%	0.5%	6.5%	8.48%
CRACK	0.8%	0.1%	0.5%	1.06%
HALLUCINOGENS	2.4%	0.3%	1.8%	2.96%
UPPERS	2.3%	0.4%	1.5%	3.11%
DOWNERS	2.5%	0.4%	1.7%	3.19%
ROHYPNOL	1.7%	0.3%	1.1%	2.20%
STEROIDS	0.5%	0.1%	0.3%	0.61%
ECSTASY	2.6%	0.4%	1.8%	3.47%
HEROIN	0.5%	0.2%	0.1%	0.94%

Lifetime Use of Selected Substances - Non-Border Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	34.8%	32.7%	36.8%	1.02%
CIGARETTES	32.9%	31.0%	34.8%	0.94%
SMOKELESS TOBACCO	10.4%	9.3%	11.5%	0.55%
ALCOHOL	65.6%	63.8%	67.4%	0.89%
INHALANTS	16.3%	15.6%	17.1%	0.39%
ANY ILLICIT DRUG	28.2%	26.6%	29.7%	0.77%
MARIJUANA	26.8%	25.3%	28.3%	0.75%
MARIJUANA ONLY	14.5%	13.6%	15.4%	0.44%
COCAINE OR CRACK	6.9%	6.1%	7.7%	0.39%
COCAINE	6.3%	5.5%	7.1%	0.39%
CRACK	2.1%	2.0%	2.3%	0.09%
HALLUCINOGENS	4.9%	4.3%	5.5%	0.29%
UPPERS	6.2%	5.4%	7.0%	0.41%
DOWNERS	6.4%	5.6%	7.2%	0.39%
ROHYPNOL	2.1%	1.8%	2.3%	0.11%
STEROIDS	1.6%	1.4%	1.7%	0.07%
ECSTASY	5.5%	5.0%	6.0%	0.25%
HEROIN	1.5%	1.4%	1.7%	0.07%

DOWNERS	0.4%	5.0%	7.270	0.59%			
ROHYPNOL	2.1%	1.8%	2.3%	0.11%			
STEROIDS	1.6%	1.4%	1.7%	0.07%			
ECSTASY	5.5%	5.0%	6.0%	0.25%			
HEROIN	1.5%	1.4%	1.7%	0.07%			
Lifetime Use of Selected Substances - Non-Border Grade 7							
CATEGORY	Prevalence	Bound	daries	Standard			
		Lower	Upper	Error			
TOBACCO	19.0%	17.0%	20.9%	0.95%			
CIGARETTES	17.9%	16.0%	19.7%	0.94%			
SMOKELESS TOBACCO	4.8%	4.2%	5.3%	0.27%			
ALCOHOL	47.0%	44.5%	49.5%	1.25%			
INHALANTS	18.4%	17.0%	19.8%	0.69%			
ANY ILLICIT DRUG	11.2%	9.4%	13.0%	0.90%			
MARIJUANA	9.7%	8.1%	11.4%	0.82%			
MARIJUANA ONLY	5.7%	4.7%	6.8%	0.51%			
COCAINE OR CRACK	3.1%	2.4%	3.7%	0.32%			
COCAINE	2.2%	1.7%	2.7%	0.26%			
CRACK	1.8%	1.5%	2.2%	0.17%			
HALLUCINOGENS	1.2%	0.9%	1.5%	0.15%			
UPPERS	0.9%	0.7%	1.2%	0.11%			

DOWNERS

ROHYPNOL

0.7%

0.7%

1.2%

1.3%

1.0%

1.0%

Past 30-day Use of Selected Substances - Non-Border Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	15.0%	13.7%	16.2%	0.63%
CIGARETTES	13.5%	12.4%	14.6%	0.55%
SMOKELESS TOBACCO	4.0%	3.5%	4.5%	0.25%
ALCOHOL	30.8%	29.2%	32.3%	0.77%
INHALANTS	5.4%	4.6%	6.1%	0.38%
ANY ILLICIT DRUG	12.7%	11.8%	13.6%	0.43%
MARIJUANA	11.3%	10.5%	12.0%	0.39%
MARIJUANA ONLY	4.5%	4.1%	5.0%	0.22%
COCAINE OR CRACK	2.4%	2.1%	2.7%	0.14%
COCAINE	2.2%	1.9%	2.4%	0.14%
CRACK	0.6%	0.5%	0.7%	0.03%
HALLUCINOGENS	1.5%	1.3%	1.7%	0.09%
UPPERS	2.5%	2.2%	2.9%	0.18%
DOWNERS	2.7%	2.3%	3.1%	0.18%
ROHYPNOL	0.8%	0.7%	0.9%	0.05%
STEROIDS	0.6%	0.5%	0.7%	0.03%
ECSTASY	1.9%	1.6%	2.1%	0.13%
HEROIN	0.5%	0.5%	0.6%	0.03%

Past 30-day Use of Selected Substances - Non-Border Grade 7				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	6.1%	5.2%	7.0%	0.46%
CIGARETTES	5.4%	4.6%	6.2%	0.40%
SMOKELESS TOBACCO	1.5%	1.2%	1.8%	0.16%
ALCOHOL	16.0%	14.2%	17.7%	0.88%
INHALANTS	7.3%	6.3%	8.4%	0.53%
ANY ILLICIT DRUG	4.8%	3.7%	5.9%	0.54%
MARIJUANA	4.0%	3.1%	4.9%	0.45%
MARIJUANA ONLY	2.0%	1.4%	2.5%	0.26%
COCAINE OR CRACK	0.9%	0.6%	1.1%	0.13%
COCAINE	0.6%	0.4%	0.8%	0.11%
CRACK	0.5%	0.4%	0.7%	0.07%
HALLUCINOGENS	0.5%	0.3%	0.6%	0.08%
UPPERS	0.5%	0.3%	0.6%	0.07%
DOWNERS	0.4%	0.3%	0.5%	0.07%
ROHYPNOL	0.5%	0.3%	0.7%	0.09%

0.10%

0.15%

STEROIDS	1.4%	1.2%	1.6%	0.08%
ECSTASY	1.6%	1.3%	1.9%	0.13%
HEROIN	1.1%	0.9%	1.3%	0.11%

STEROIDS	0.5%	0.4%	0.6%	0.05%
ECSTASY	0.5%	0.4%	0.6%	0.07%
HEROIN	0.4%	0.3%	0.6%	0.06%

Lifetime Use of Selected Substances - Non-Border Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	26.8%	24.6%	29.1%	1.13%
CIGARETTES	25.1%	22.9%	27.3%	1.10%
SMOKELESS TOBACCO	7.2%	6.2%	8.2%	0.48%
ALCOHOL	59.3%	57.0%	61.7%	1.18%
INHALANTS	19.9%	18.7%	21.1%	0.60%
ANY ILLICIT DRUG	18.3%	16.2%	20.3%	1.01%
MARIJUANA	16.8%	14.9%	18.8%	0.99%
MARIJUANA ONLY	9.9%	8.8%	11.1%	0.59%
COCAINE OR CRACK	4.1%	3.4%	4.7%	0.32%
COCAINE	3.2%	2.7%	3.8%	0.28%
CRACK	2.0%	1.7%	2.3%	0.15%
HALLUCINOGENS	2.5%	2.1%	2.9%	0.18%
UPPERS	2.6%	2.2%	3.0%	0.19%
DOWNERS	2.4%	1.9%	2.8%	0.21%
ROHYPNOL	1.8%	1.4%	2.2%	0.18%
STEROIDS	1.5%	1.3%	1.8%	0.13%
ECSTASY	3.1%	2.7%	3.5%	0.19%
HEROIN	1.7%	1.4%	1.9%	0.11%

Past 30-day Use of Selected Substances - Non-Border Grade 8				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	8.8%	7.7%	9.9%	0.55%
CIGARETTES	7.9%	6.9%	8.9%	0.50%
SMOKELESS TOBACCO	2.3%	1.9%	2.8%	0.23%
ALCOHOL	22.3%	20.4%	24.1%	0.92%
INHALANTS	7.3%	6.0%	8.5%	0.61%
ANY ILLICIT DRUG	7.6%	6.6%	8.7%	0.52%
MARIJUANA	6.7%	5.7%	7.7%	0.50%
MARIJUANA ONLY	3.2%	2.7%	3.7%	0.27%
COCAINE OR CRACK	1.4%	1.2%	1.6%	0.11%
COCAINE	1.1%	0.9%	1.3%	0.11%
CRACK	0.6%	0.5%	0.7%	0.05%
HALLUCINOGENS	0.7%	0.6%	0.8%	0.06%
UPPERS	1.0%	0.8%	1.2%	0.10%
DOWNERS	0.9%	0.7%	1.1%	0.12%
ROHYPNOL	0.7%	0.5%	0.8%	0.08%
STEROIDS	0.5%	0.4%	0.6%	0.06%
ECSTASY	1.0%	0.8%	1.2%	0.11%
HEROIN	0.5%	0.4%	0.6%	0.05%

Lifetime Use of Selected Substances - Non-Border Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	35.2%	33.0%	37.4%	1.10%
CIGARETTES	33.0%	30.9%	35.1%	1.06%
SMOKELESS TOBACCO	9.9%	8.7%	11.1%	0.61%
ALCOHOL	67.1%	65.4%	68.8%	0.87%
INHALANTS	17.5%	16.2%	18.8%	0.67%
ANY ILLICIT DRUG	27.8%	25.9%	29.6%	0.92%
MARIJUANA	26.1%	24.3%	28.0%	0.93%
MARIJUANA ONLY	14.6%	13.3%	16.0%	0.69%
COCAINE OR CRACK	6.2%	5.5%	6.9%	0.36%
COCAINE	5.5%	4.8%	6.2%	0.35%
CRACK	2.3%	1.9%	2.6%	0.17%
HALLUCINOGENS	4.5%	3.9%	5.1%	0.31%

Past 30-day Use of Selected Substances - Non-Border Grade 9					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	14.0%	13.0%	15.1%	0.52%	
CIGARETTES	12.4%	11.4%	13.3%	0.46%	
SMOKELESS TOBACCO	3.8%	3.2%	4.4%	0.29%	
ALCOHOL	30.6%	29.1%	32.1%	0.75%	
INHALANTS	6.0%	4.9%	7.0%	0.52%	
ANY ILLICIT DRUG	12.6%	11.5%	13.6%	0.51%	
MARIJUANA	11.0%	9.9%	12.1%	0.54%	
MARIJUANA ONLY	4.9%	4.1%	5.6%	0.38%	
COCAINE OR CRACK	2.3%	2.0%	2.7%	0.18%	
COCAINE	2.1%	1.7%	2.4%	0.16%	
CRACK	0.7%	0.5%	0.9%	0.09%	
HALLUCINOGENS	1.4%	1.2%	1.7%	0.12%	

_				
UPPERS	5.3%	4.6%	6.0%	0.35%
DOWNERS	5.4%	4.7%	6.1%	0.34%
ROHYPNOL	2.4%	2.1%	2.7%	0.16%
STEROIDS	1.4%	1.2%	1.6%	0.11%
ECSTASY	5.3%	4.8%	5.8%	0.25%
HEROIN	1.8%	1.5%	2.0%	0.14%

UPPERS	2.0%	1.7%	2.4%	0.17%
DOWNERS	2.2%	1.8%	2.6%	0.19%
ROHYPNOL	0.9%	0.7%	1.1%	0.10%
STEROIDS	0.5%	0.4%	0.6%	0.05%
ECSTASY	2.0%	1.6%	2.3%	0.16%
HEROIN	0.6%	0.4%	0.8%	0.09%

Lifetime Use of Sele	Lifetime Use of Selected Substances - Non-Border Grade 10				
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	39.7%	37.5%	41.8%	1.08%	
CIGARETTES	37.4%	35.3%	39.4%	1.02%	
SMOKELESS TOBACCO	12.2%	10.9%	13.5%	0.64%	
ALCOHOL	72.3%	70.5%	74.1%	0.91%	
INHALANTS	15.8%	14.6%	17.0%	0.60%	
ANY ILLICIT DRUG	34.2%	32.2%	36.2%	0.99%	
MARIJUANA	32.8%	30.9%	34.7%	0.95%	
MARIJUANA ONLY	17.2%	15.9%	18.5%	0.63%	
COCAINE OR CRACK	8.0%	7.0%	9.1%	0.51%	
COCAINE	7.6%	6.6%	8.6%	0.50%	
CRACK	2.2%	1.9%	2.5%	0.16%	
HALLUCINOGENS	6.8%	6.1%	7.5%	0.37%	
UPPERS	8.4%	7.3%	9.4%	0.51%	
DOWNERS	8.7%	7.7%	9.7%	0.50%	
ROHYPNOL	2.7%	2.3%	3.1%	0.19%	
STEROIDS	1.8%	1.6%	2.1%	0.13%	
ECSTASY	7.0%	6.2%	7.7%	0.36%	
HEROIN	1.5%	1.2%	1.7%	0.15%	

Past 30-day Use of Selected Substances - Non-Border Grade 10					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.5%	16.0%	18.9%	0.72%	
CIGARETTES	15.7%	14.4%	17.0%	0.67%	
SMOKELESS TOBACCO	4.7%	4.2%	5.2%	0.26%	
ALCOHOL	36.1%	34.5%	37.7%	0.80%	
INHALANTS	4.8%	3.7%	5.9%	0.54%	
ANY ILLICIT DRUG	15.8%	14.8%	16.9%	0.52%	
MARIJUANA	13.7%	12.8%	14.7%	0.48%	
MARIJUANA ONLY	5.2%	4.7%	5.7%	0.25%	
COCAINE OR CRACK	2.7%	2.3%	3.1%	0.21%	
COCAINE	2.5%	2.1%	2.9%	0.21%	
CRACK	0.6%	0.5%	0.7%	0.06%	
HALLUCINOGENS	2.1%	1.7%	2.5%	0.20%	
UPPERS	3.8%	3.3%	4.3%	0.27%	
DOWNERS	4.1%	3.6%	4.7%	0.28%	
ROHYPNOL	0.9%	0.7%	1.1%	0.09%	
STEROIDS	0.6%	0.5%	0.8%	0.06%	
ECSTASY	2.5%	2.0%	2.9%	0.21%	
HEROIN	0.6%	0.5%	0.7%	0.07%	

Lifetime Use of Selected Substances - Non-Border Grade 11				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	43.9%	41.7%	46.1%	1.10%
CIGARETTES	41.8%	39.8%	43.8%	1.00%
SMOKELESS TOBACCO	14.3%	12.9%	15.8%	0.74%
ALCOHOL	74.6%	73.0%	76.2%	0.79%
INHALANTS	13.1%	11.8%	14.3%	0.62%
ANY ILLICIT DRUG	40.1%	38.5%	41.6%	0.76%
MARIJUANA	38.8%	37.2%	40.3%	0.77%
MARIJUANA ONLY	20.3%	19.1%	21.5%	0.59%
COCAINE OR CRACK	10.4%	9.2%	11.6%	0.61%

Past 30-day Use of Selected Substances - Non-Border Grade 11				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	21.6%	19.8%	23.3%	0.86%
CIGARETTES	19.7%	18.1%	21.2%	0.78%
SMOKELESS TOBACCO	5.9%	5.3%	6.5%	0.30%
ALCOHOL	39.4%	37.6%	41.2%	0.89%
INHALANTS	3.5%	2.8%	4.2%	0.36%
ANY ILLICIT DRUG	18.6%	17.3%	20.0%	0.68%
MARIJUANA	16.8%	15.5%	18.0%	0.64%
MARIJUANA ONLY	6.3%	5.5%	7.0%	0.39%
COCAINE OR CRACK	3.8%	3.3%	4.3%	0.26%

COCAINE	10.1%	8.9%	11.3%	0.61%
CRACK	2.3%	2.0%	2.7%	0.18%
HALLUCINOGENS	7.3%	6.1%	8.4%	0.58%
UPPERS	10.7%	9.2%	12.2%	0.76%
DOWNERS	11.3%	9.9%	12.7%	0.71%
ROHYPNOL	2.4%	1.9%	2.9%	0.24%
STEROIDS	1.6%	1.3%	1.8%	0.12%
ECSTASY	8.1%	6.9%	9.2%	0.56%
HEROIN	1.7%	1.4%	2.0%	0.14%

COCAINE	3.6%	3.1%	4.1%	0.26%
CRACK	0.6%	0.4%	0.8%	0.09%
HALLUCINOGENS	2.1%	1.6%	2.6%	0.25%
UPPERS	4.5%	3.7%	5.3%	0.39%
DOWNERS	4.8%	4.1%	5.6%	0.37%
ROHYPNOL	0.9%	0.7%	1.2%	0.12%
STEROIDS	0.7%	0.5%	0.8%	0.07%
ECSTASY	2.6%	2.0%	3.2%	0.30%
HEROIN	0.4%	0.3%	0.6%	0.07%

Lifetime Use of Selected Substances - Non-Border Grade 12					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	48.7%	45.6%	51.8%	1.53%	
CIGARETTES	46.7%	43.7%	49.8%	1.51%	
SMOKELESS TOBACCO	16.2%	14.1%	18.2%	1.03%	
ALCOHOL	77.4%	75.5%	79.4%	0.98%	
INHALANTS	11.6%	10.6%	12.6%	0.49%	
ANY ILLICIT DRUG	42.9%	40.7%	45.1%	1.08%	
MARIJUANA	41.9%	39.7%	44.0%	1.06%	
MARIJUANA ONLY	21.7%	20.6%	22.8%	0.55%	
COCAINE OR CRACK	11.3%	9.7%	12.9%	0.79%	
COCAINE	11.0%	9.4%	12.6%	0.79%	
CRACK	2.2%	1.9%	2.6%	0.19%	
HALLUCINOGENS	8.5%	7.4%	9.4%	0.50%	
UPPERS	11.2%	9.5%	12.9%	0.83%	
DOWNERS	12.0%	10.5%	13.5%	0.75%	
ROHYPNOL	2.1%	1.7%	2.5%	0.21%	
STEROIDS	1.8%	1.5%	2.1%	0.15%	
ECSTASY	9.1%	8.0%	10.2%	0.53%	
HEROIN	1.5%	1.2%	1.7%	0.13%	

Past 30-day Use of Selected Substances - Non-Border Grade 12				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	25.3%	22.8%	27.8%	1.25%
CIGARETTES	23.6%	21.3%	25.8%	1.12%
SMOKELESS TOBACCO	7.0%	5.6%	8.3%	0.66%
ALCOHOL	45.2%	42.5%	48.0%	1.38%
INHALANTS	2.4%	1.9%	2.8%	0.24%
ANY ILLICIT DRUG	19.3%	17.9%	20.8%	0.72%
MARIJUANA	17.6%	16.4%	18.9%	0.63%
MARIJUANA ONLY	6.5%	5.8%	7.1%	0.31%
COCAINE OR CRACK	3.8%	3.2%	4.4%	0.30%
COCAINE	3.6%	3.0%	4.2%	0.30%
CRACK	0.5%	0.4%	0.7%	0.08%
HALLUCINOGENS	2.4%	2.0%	2.7%	0.17%
UPPERS	4.2%	3.4%	5.0%	0.39%
DOWNERS	4.7%	3.8%	5.5%	0.40%
ROHYPNOL	0.7%	0.6%	0.9%	0.08%
STEROIDS	0.8%	0.6%	1.0%	0.10%
ECSTASY	2.9%	2.3%	3.6%	0.31%
HEROIN	0.5%	0.3%	0.7%	0.08%